

Geography of local configurations

D. Coupier

February 1, 2008

Université Lille 1

E-Mail address : david.coupier@math.univ-lille1.fr

Mail address : Laboratoire Paul Painlevé, Université Lille 1
Cité scientifique, 59655 Villeneuve d'Ascq Cedex, France.

Telephone : 33 (0)3 20 43 67 60

Fax : 33 (0)3 20 43 43 02

Abstract

A d -dimensional ferromagnetic Ising model on a lattice torus is considered. As the size n of the lattice tends to infinity, the magnetic field $a = a(n)$ and the pair potential depend on n . Precise bounds for the probability for local configurations to occur in a large ball are given. Under some conditions bearing on potentials $a(n)$ and $b(n)$, the distance between copies of different local configurations is estimated according to their weights. Finally, a sufficient condition ensuring that a given local configuration occurs everywhere in the lattice is suggested.

Key words : Ising model, ferromagnetic interaction, FKG inequality.

AMS Subject Classification : 60F05, 82B20.

1 Introduction

In the theory of random graphs, inaugurated by Erdős and Rényi [12], the appearance of a given subgraph has been widely studied (see [4] or [22] for a general reference). In the random graph formed by n vertices, in which the edges are chosen independently with probability $0 < p < 1$, a subgraph may occur or not according to the value of $p = p(n)$. In addition, under a certain condition on the probability $p(n)$, its number of occurrences in the graph is asymptotically (i.e. as $n \rightarrow +\infty$) Poissonian. Replacing the edges with the spins of an Ising model, the notion of subgraph corresponds to the notion of what we will call *local configuration*; Figure 1 shows an example. Using tools coming from random graphs (as *threshold functions* and *Poisson approximations*), the study of the appearance of a given local configuration has been done in [11], [10] and [8]. In this article, this study is extended into three directions. First, the speed at which local configurations occur is precised. Moreover, when the number of copies in the graph of a given local configuration is finite, the geography of positive and negative spins surrounding one of them is described. Finally, a sufficient condition ensuring that a given local configuration is present everywhere in the graph is stated. The results obtained in these three directions are based on the same tools; the Markovian character of the measure, the control of the conditional probability for a local configuration to occur in the graph and the FKG inequality [15].

Let us consider a lattice graph in dimension $d \geq 1$, with periodic boundary conditions (lattice torus). The vertex set is $V_n = \{0, \dots, n-1\}^d$. The integer n will be called the *size* of the lattice. The edge set, denoted by E_n , will be specified by defining the set of neighbors $\mathcal{V}(x)$ of a given vertex x :

$$\mathcal{V}(x) = \{y \neq x \in V_n, \|y - x\|_q \leq \rho\}, \quad (1)$$

where the subtraction is taken componentwise modulo n , $\|\cdot\|_q$ stands for the L_q norm in $\mathbb{R}^d$ ($1 \leq q \leq \infty$), and ρ is a fixed integer. For instance, the square lattice is obtained for $q = \rho = 1$. Replacing the L_1 norm with the L_∞ norm adds the diagonals. From now on, all operations on vertices will be understood modulo n . In particular, each vertex of the lattice has the same number of neighbors; we denote by $\mathcal{V}$ this number.

A *configuration* is a mapping from the vertex set V_n to the state space $\{-1, +1\}$. Their set is denoted by $\mathcal{X}_n = \{-1, +1\}^{V_n}$ and called the *configuration set*. The Ising model is classically defined as follows (see e.g. Georgii [17] and Malyshev and Minlos [21]).

Definition 1.1 *Let $G_n = (V_n, E_n)$ be the undirected graph structure with finite vertex set V_n and edge set E_n . Let a and b be two reals. The Ising model with parameters a and b is the probability measure $\mu_{a,b}$ on $\mathcal{X}_n = \{-1, +1\}^{V_n}$ defined by: for all $\sigma \in \mathcal{X}_n$,*

$$\mu_{a,b}(\sigma) = \frac{1}{Z_{a,b}} \exp \left(a \sum_{x \in V_n} \sigma(x) + b \sum_{\{x,y\} \in E_n} \sigma(x)\sigma(y) \right), \quad (2)$$

where the normalizing constant $Z_{a,b}$ is such that $\sum_{\sigma \in \mathcal{X}_n} \mu_{a,b}(\sigma) = 1$.

Following the definition of [21] p. 2, the measure $\mu_{a,b}$ defined above is a Gibbs measure associated to potentials a and b . Expectations relative to $\mu_{a,b}$ will be denoted by $\mathbb{E}_{a,b}$. In the classical presentation of statistical physics, the elements of $\mathcal{X}_n$ are spin configurations; each vertex of V_n is an atom whose spin is either positive or negative. Here, we shall simply talk about positive or negative vertices instead of positive or negative spins and we shall merely denote by $+$ and $-$ the states $+1$ and -1 . The parameters a and b are respectively the *magnetic field* and the *pair potential*. The model remaining unchanged by swapping positive and negative vertices and replacing a by $-a$, we chose to study only negative values of the magnetic field a . Throughout the paper, in order to use the FKG inequality, the pair potential b will be supposed nonnegative.

As the size n of the lattice tends to infinity, the potentials $a = a(n)$ and $b = b(n)$ depend on n . The case where $a(n)$ tends to $-\infty$ corresponds to rare positive vertices among a majority of negative ones. In order to simplify formulas, the Gibbs measure $\mu_{a(n),b(n)}$ will be merely denoted by $\mu_{a,b}$.

We are interested in the appearance in the graph G_n of families of local configurations. See Section 2 for a precise definition and Figure 1 for an example. Such configurations are called “local” in the sense that the vertex set on which they are defined is fixed and does not depend on n . A local configuration η is determined by its set of positive vertices $V_+(\eta)$ whose cardinality and perimeter are respectively denoted by $k(\eta)$ and $\gamma(\eta)$. A natural idea (coming from [8]) consists in regarding both parameters $k(\eta)$ and $\gamma(\eta)$ through the same quantity; the *weight* of the local configuration η

$$W_n(\eta) = \exp(2a(n)k(\eta) - 2b(n)\gamma(\eta)) \ .$$

This notion plays a central role in our study. Indeed, the weight $W_n(\eta)$ represents the probabilistic cost associated to a given occurrence of η .




Figure 1: A local configuration η with $k(\eta) = |V_+(\eta)| = 10$ positive vertices and a perimeter $\gamma(\eta)$ equals to 58, in dimension $d = 2$ and on a ball of radius $r = 2$ (with $\rho = 1$ and relative to the L_∞ norm).

Proving some sharp inequalities is generally more difficult than stating only limits. In the case of random graphs, Janson et al. [20], thus Janson [19], have obtained exponential bounds for the probability of nonexistence of subgraphs. Some other useful inequalities have been suggested by Boppona and Spencer [5]. In bond percolation on $\mathbb{Z}^d$, it is believed that, in the subcritical phase, the probability for the radius of an open cluster of being

larger than n behaves as an exponential term multiplied by a power of n : see [18] p. 85 for precise bounds. But, when the variables of the system are dependent, as the spins of the Ising model, such inequalities become harder to obtain. The Stein-Chen method (see Barbour et al. [3] for a very complete reference or [7] for the original paper of Chen) is a useful way to bound the error of a Poisson approximation and so, in particular, to bound the absolute value of the difference between the probability of a property of the model and its limit. An example of such a property is the appearance of a negative vertex (see Ganesh et al. [16]) or more generally that of any given local configuration [10]. These two previous papers concern the case of a divergent magnetic field $|a|$ and a constant pair potential. Coupling with the loss-network space-time representation due to Fernández et al. [13], Ferrari and Picco [14] have proved an exponential bound for large contours at low temperature (i.e. b large enough) and zero magnetic field. Finally, under mixing conditions, various exponential approximations with error bounds have been proved; see Abadi and Galves [2] for an overview and Abadi et al. [1] for the high temperature case (i.e. b small enough).

Our first goal is to establish precise lower and upper bounds for the probability for certain families of local configurations to occur in the graph for unbounded potentials $a(n)$ and $b(n)$.

Let $x \in V_n$ be a vertex, $W \in]0, 1[$ be a real number and R be an integer. We denote by $\mathcal{A}(x, R, W)$ the (interpreted) event “a local configuration whose weight is smaller than W occurs somewhere in the ball of center x and radius R ”. The study of the probability of this event becomes interesting when the radius $R = R(n)$ and the weight $W = W_n$ depend on n and tend respectively to infinity and zero. Then, Theorem 4.1 gives precise bounds for the probability of $\mathcal{A}(x, R(n), W_n)^c$. There exist two (explicit) constants $K > K' > 0$ such that for n large enough,

$$e^{-KR(n)^d W_n} \leq 1 - \mu_{a,b}(\mathcal{A}(x, R(n), W_n)) \leq e^{-K'R(n)^d W_n}.$$

Another interesting problem consists in describing geographically what happens in the studied model: the size of objects occurring in the model and the distance between them. The size of components in random graphs or the radius of open clusters in percolation are two classical examples (see respectively [4] and [18]). For the low-temperature plus-phase of the Ising model, Chazottes and Redig [6] have studied the appearance of the first two copies of a given pattern in terms of occurrence time and repetition time. The occurrence time T_A of a pattern A represents the volume of the smallest set of vertices in which A can be found. As the size of the pattern increases, the distribution of T_A is approximated by an exponential law with error bounds. The same is true for the repetition time R_A . Similar results exist for sufficiently mixing Gibbs random fields (see [1]).

In our context, the studied objects are local. Hence, our second goal is to estimate the distance between different copies of local configurations occurring in the graph.

Let η be a local configuration. It has been proved in [8] that the number of copies of η occurring in G_n is asymptotically Poissonian provided the product $n^d W_n(\eta)$ is constant (the precise hypotheses are denoted by (H) and recalled in the beginning of Section 5). In particular, the number of copies of η in the graph is finite with probability tending to 1. Let

η_x be one of them (that occurring on the ball centered at x). We first observe that vertices surrounding η_x are all negative (Lemma 5.1). Hence, a natural question is the distance from η_x to the closest positive vertices. Theorem 5.3 answers to this question; it states that the distance from η_x to the closest local configurations of weight W_n is of order $W_n^{-1/d}$. Two other results complete the study of the geography of local configurations under the hypothesis (H) . Theorem 5.4 claims the distance between the closest local configurations (to η_x) of weight W_n is also of order $W_n^{-1/d}$. The situation described by Theorems 5.3 and 5.4 is represented in Figure 4. Finally, Theorem 5.3 implies the distance between two any copies of η should be of order n , which is the size of the graph. Proposition 5.5 precises this intuition.

From [8], a condition ensuring that a local configuration η occurs in G_n is deduced; if the product $n^d W_n(\eta)$ tends to infinity then, with probability tending to 1, at least one copy of η can be found somewhere in the graph. However, an uncertainty remains about the places in G_n where η occurs. A richer information would be to know when the local configuration η occurs everywhere in G_n . Inequalities stated in the proof of Theorem 4.1 allow us to obtain such an information.

For that purpose, the lattice V_n is divided into blocks of $(2R(n) + 1)^d$ vertices. Thus, a supergraph $\tilde{G}_n$ whose set of vertices $\tilde{V}_n$ is formed by the centers of these blocks is defined. In order to study the appearance of η in each of these blocks, the set of configurations of $\tilde{V}_n$ is endowed with an appropriate measure $\tilde{\mu}_{a,b}$ (depending on η). Proposition 6.2 precises the asymptotic behavior of $\tilde{\mu}_{a,b}$ according to the weight $W_n(\eta)$, the radius of the blocks $R(n)$ and the size n . In particular, if

$$\lim_{n \rightarrow +\infty} R(n)^d \ln \left(\frac{n}{R(n)} \right)^{-1} W_n(\eta) = +\infty$$

then, with probability tending to 1, all the blocks contain at least one copy of the local configuration η .

The paper is organized as follows. The notion of local configuration η is defined in Section 2. Its number of positive vertices $k(\eta)$, its perimeter $\gamma(\eta)$ and its weight $W_n(\eta)$ are also introduced. Section 3 is devoted to the three main tools of our study. Property 3.1 underlines the Markovian character of the Gibbs measure $\mu_{a,b}$. Thus, a control of the conditional probability for a local configuration to occur on a ball uniformly on the neighborhood of that ball is given in Lemma 3.2. Finally, the FKG inequality is discussed at the end of Section 3. Section 4 gives the proof of Theorem 4.1. The geography of local configurations occurring in the Ising model is described in Section 5. Section 5.1 introduces the problem (the hypothesis (H) and Lemma 5.1). Theorems 5.3 and 5.4 and Proposition 5.5 are stated in Section 5.2 and proved in Section 5.3. The graph $\tilde{G}_n$ and the measure $\tilde{\mu}_{a,b}$ are defined in Section 6. The latter inherits from $\mu_{a,b}$ its Markovian character (Property 6.1). Finally, Proposition 6.2 studies its asymptotic behavior.

2 Local configurations

Let us start with some notations and definitions. Given $\zeta \in \mathcal{X}_n = \{-, +\}^{V_n}$ and $V \subset V_n$, we denote by ζ_V the natural projection of ζ over $\{-, +\}^V$. If U and V are two disjoint subsets of V_n then $\zeta_U \zeta'_V$ is the configuration on $U \cup V$ which is equal to ζ on U and ζ' on V . Let us denote by δV the neighborhood of V (corresponding to (1)):

$$\delta V = \{y \in V_n \setminus V, \exists x \in V, \{x, y\} \in E_n\} ,$$

and by $\overline{V}$ the union of the two disjoint sets V and δV . Moreover, $|V|$ denotes the cardinality of V and $\mathcal{F}(V)$ the σ -algebra generated by the configurations of $\{-, +\}^V$.

As usual, the graph distance $dist$ is defined as the minimal length of a path between two vertices. We shall denote by $B(x, r)$ the ball of center x and radius r :

$$B(x, r) = \{y \in V_n; dist(x, y) \leq r\} .$$

In the case of balls, $\overline{B(x, r)} = B(x, r+1)$. In order to avoid unpleasant situations, like self-overlapping balls, we will always assume that $n > 2\rho r$. If n and n' are both larger than $2\rho r$, the balls $B(x, r)$ in G_n and $G_{n'}$ are isomorphic. Two properties of the balls $B(x, r)$ will be crucial in what follows. The first one is that two balls with the same radius are translates of each other:

$$B(x + y, r) = y + B(x, r) .$$

The second one is that for $n > 2\rho r$, the cardinality of $B(x, r)$ depends only on r and neither on x nor on n : it will be denoted by β_r .

Let r be a positive integer, and consider a fixed ball with radius r , say $B(0, r)$. We denote by $\mathcal{C}_r = \{-, +\}^{B(0, r)}$ the set of configurations on that ball. Elements of $\mathcal{C}_r$ will be called *local configurations with radius r* , or merely *local configurations* whenever the radius r will be fixed. Of course, there exists only a finite number of such configurations (precisely 2^{β_r}). See Figure 1 for an example. Throughout this paper, the radius r will be constant, i.e. it will not depend on the size n . Hence, defining local configurations on balls of radius r serves only to ensure that studied objects are “local”. In what follows, η, η' will denote local configurations of radius r .

A local configuration $\eta \in \mathcal{C}_r$ is determined by its subset $V_+(\eta) \subset B(0, r)$ of positive vertices:

$$V_+(\eta) = \{x \in B(0, r), \eta(x) = +\} .$$

The cardinality of this set will be denoted by $k(\eta)$ and its complement in $B(0, r)$, i.e. the set of negative vertices of η , by $V_-(\eta)$. Moreover, the geometry (in the sense of the graph structure) of the set $V_+(\eta)$ needs to be described. Let us define the *perimeter* $\gamma(\eta)$ of the local configuration η by the formula:

$$\gamma(\eta) = \mathcal{V} |V_+(\eta)| - 2 |\{\{x, y\} \in V_+(\eta) \times V_+(\eta), \{x, y\} \in E_n\}| ,$$

where $\mathcal{V}$ is the number of neighbors of a vertex. In other words, $\gamma(\eta)$ counts the pairs of neighboring vertices x and y of $B(0, r)$ having opposite spins (under η) and those such that $x \in B(0, r)$, $y \in \delta B(0, r)$ and $\eta(x) = +$.

We denote by $W_n(\eta)$ and call the *weight* of the local configuration η the following quantity:

$$W_n(\eta) = \exp(2a(n)k(\eta) - 2b(n)\gamma(\eta)) .$$

Since $a(n) < 0$ and $b(n) \geq 0$, the weight $W_n(\eta)$ satisfies $0 < W_n(\eta) \leq 1$. That of the local configuration having only negative vertices, denoted by η^0 and called the *null local configuration*, is equal to 1. If $\eta \neq \eta^0$ then $k(\eta) \geq 1$ and $\gamma(\eta) \geq \mathcal{V}$. It follows that

$$W_n(\eta) \leq \exp(2a(n) - 2b(n)\mathcal{V}) .$$

Actually, the weight $W_n(\eta)$ represents the probabilistic cost associated to the presence of η on a given ball. This idea will be precised in the next section.

Remark the notations $k(\cdot)$, $\gamma(\cdot)$ and $W_n(\cdot)$ can be naturally extended to any configuration $\zeta \in \{-, +\}^V$, $V \subset V_n$.

Let $\eta \in \mathcal{C}_r$. For each vertex $x \in V_n$, denote by η_x the translation of η onto the ball $B(x, r)$ (up to periodic boundary conditions):

$$\forall y \in V_n, \text{ dist}(0, y) \leq r \implies \eta_x(x + y) = \eta(y) .$$

In particular, $V_+(\eta_x) = x + V_+(\eta)$. So, η and η_x have the same number of positive vertices and the same perimeter. So do their weights. Let us denote by I_x^η the indicator function defined on $\mathcal{X}_n$ as follows: $I_x^\eta(\sigma)$ is 1 if the restriction of the configuration $\sigma \in \mathcal{X}_n$ to the ball $B(x, r)$ is η_x and 0 otherwise.

Let V and V' be two disjoint subsets of vertices. The following relations

$$k(\zeta\zeta') = k(\zeta) + k(\zeta') \quad \text{and} \quad \gamma(\zeta\zeta') \leq \gamma(\zeta) + \gamma(\zeta')$$

are true whatever the configurations $\zeta \in \{-, +\}^V$ and $\zeta' \in \{-, +\}^{V'}$. As an immediate consequence, the weight $W_n(\zeta\zeta')$ is larger than the product $W_n(\zeta)W_n(\zeta')$. The *connection* between $\zeta \in \{-, +\}^V$ and $\zeta' \in \{-, +\}^{V'}$, denoted by $\text{conn}(\zeta, \zeta')$, is defined by:

$$\text{conn}(\zeta, \zeta') = |\{\{y, z\} \in E_n, y \in V, z \in V' \text{ and } \zeta(y) = \zeta'(z) = +\}| .$$

This quantity allows to link the perimeters of the configurations $\zeta\zeta'$, ζ and ζ' together:

$$\gamma(\zeta\zeta') + 2\text{conn}(\zeta, \zeta') = \gamma(\zeta) + \gamma(\zeta') ,$$

and therefore their weights:

$$W_n(\zeta\zeta') \exp(-4b(n)\text{conn}(\zeta, \zeta')) = W_n(\zeta)W_n(\zeta') . \quad (3)$$

In particular, if the connection $\text{conn}(\zeta, \zeta')$ is null then the weight $W_n(\zeta\zeta')$ is equal to the product $W_n(\zeta)W_n(\zeta')$. This is the case when $\overline{V} \cap V' = \emptyset$.

3 The three main tools

This section is devoted to the main tools on which are based all the results of this paper: the Markovian character of the Gibbs measure $\mu_{a,b}$, a control of the probability for $\eta \in \mathcal{C}_r$ to occur on a given ball and the FKG inequality. Excepted the first part of Lemma 3.2, the results of this section are not new.

Two subsets of vertices U and V of V_n are said $\mathcal{V}$ -disjoint if none of the vertices of U belong to the neighborhood of one of the vertices of V . In other words, U and V are $\mathcal{V}$ -disjoint if and only if $U \cap \overline{V} = \emptyset$ (or equivalently $\overline{U} \cap V = \emptyset$). For example, two balls $B(x, r)$ and $B(x', r)$ are $\mathcal{V}$ -disjoint if and only if the distance between their centers x and x' is larger than $2r + 1$, i.e $\text{dist}(x, x') > 2r + 1$.

The following result is a classical property of Gibbs measures (see [17] p.157); it describes the Markovian character of $\mu_{a,b}$. The second part of Property 3.1 means that, given two $\mathcal{V}$ -disjoint sets U and V , the σ -algebras $\mathcal{F}(U)$ and $\mathcal{F}(V)$ are conditionally independent knowing the configuration on $\delta U \cup \delta V$.

For any sets of vertices V, V' and for any event $A \in \mathcal{F}(V)$, the function $\mu_{a,b}(A|\mathcal{F}(V'))$ denotes the $\mathcal{F}(V')$ -measurable random variable defined as follows; for $\sigma \in \{-, +\}^{V'}$, $\mu_{a,b}(A|\mathcal{F}(V'))(\sigma)$ is the conditional probability $\mu_{a,b}(A|\sigma)$.

Property 3.1 1. Let $V, V' \subset V_n$ be two sets of vertices such that $V \cap V' = \emptyset$ and $\delta V \subset V'$. Then, for all $A \in \mathcal{F}(V)$,

$$\mu_{a,b}(A|\mathcal{F}(V')) = \mu_{a,b}(A|\mathcal{F}(\delta V)) . \quad (4)$$

2. Let $U, V, V' \subset V_n$ be three sets of vertices such that U and V are $\mathcal{V}$ -disjoint, $(U \cup V) \cap V' = \emptyset$ and $\delta U \cup \delta V \subset V'$. Then, for all $A \in \mathcal{F}(U)$ and $B \in \mathcal{F}(V)$,

$$\mu_{a,b}(A \cap B|\mathcal{F}(V')) = \mu_{a,b}(A|\mathcal{F}(V')) \mu_{a,b}(B|\mathcal{F}(V')) . \quad (5)$$

Let us note that (4) is a consequence of the identity

$$\forall \sigma \in \{-, +\}^{\delta V}, \forall \sigma' \in \{-, +\}^{V' \setminus \delta V}, \mu_{a,b}(A|\sigma\sigma') = \mu_{a,b}(A|\sigma) ,$$

which itself relies on the exponential form of the Gibbs measure $\mu_{a,b}$ (see Definition 1.1). The proof of a similar identity is available in [21] p.7.

Finally, remark that the second part of Property 3.1 can be easily extended to any finite family of sets of vertices which are two by two $\mathcal{V}$ -disjoint.

Let η be a local configuration. Thanks to the translation invariance of the graph G_n , the indicator functions I_x^η , $x \in V_n$, have the same distribution. So, let us pick a vertex x . For any configuration $\sigma \in \{-, +\}^{\delta B(x, r)}$, the quantity $\mu_{a,b}(I_x^\eta = 1|\sigma)$ represents the probability for η (or η_x) to occur on the ball $B(x, r)$ knowing what happens on its neighborhood. A precise study of this conditional probability have been done in [8], Section 2. This study provides lower and upper bounds for the probability for η to occur on the ball $B(x, r)$:

Lemma 3.2 *Let η be a local configuration with radius r and x be a vertex.*

On the one hand, if $a(n) + \mathcal{V}b(n) \leq 0$ then there exists a constant $c_r > 0$ such that, for all configuration $\sigma \in \{-, +\}^{\delta B(x, r)}$,

$$\mu_{a,b}(I_x^\eta = 1 | \sigma) \geq c_r W_n(\eta) . \quad (6)$$

In particular, $\mu_{a,b}(I_x^\eta = 1)$ satisfies the same lower bound. On the other hand, if $a(n) + 2\mathcal{V}b(n) \leq 0$ then there exists a constant $C_r > 0$ such that

$$\mu_{a,b}(I_x^\eta = 1) \leq C_r W_n(\eta) . \quad (7)$$

The constants C_r and c_r depend only on the radius r (and also on parameters q, ρ and d) but not on the local configuration η .

Proof: Let η be a local configuration with radius r , x be a vertex and σ be an element of $\{-, +\}^{\delta B(x, r)}$. Lemma 2.2 of [8] allows to write the conditional probability $\mu_{a,b}(I_x^\eta = 1 | \sigma)$ as a function of weights of some configurations:

$$\begin{aligned} \mu_{a,b}(I_x^\eta = 1 | \sigma) &= \frac{W_n(\eta_x \sigma)}{\sum_{\eta' \in \mathcal{C}_r} W_n(\eta'_x \sigma)} \\ &\geq W_n(\eta) \left(\sum_{\eta' \in \mathcal{C}_r} \frac{W_n(\eta'_x \sigma)}{W_n(\sigma)} \right)^{-1} , \end{aligned} \quad (8)$$

using the inequality $W_n(\eta_x \sigma) \geq W_n(\eta) W_n(\sigma)$. Let η' be a local configuration. Its perimeter $\gamma(\eta')$ can be bounded as follows:

$$\mathcal{V}k(\eta') \geq \gamma(\eta') \geq \text{conn}(\eta'_x, \sigma) .$$

Hence, $\gamma(\eta') + \mathcal{V}k(\eta')$ is larger than $2\text{conn}(\eta'_x, \sigma)$ and the difference $\gamma(\eta'_x \sigma) - \gamma(\sigma)$ satisfies:

$$\begin{aligned} \gamma(\eta'_x \sigma) - \gamma(\sigma) &= \gamma(\eta') - 2\text{conn}(\eta'_x, \sigma) \\ &\geq -\mathcal{V}k(\eta') . \end{aligned}$$

We deduce from this last inequality that

$$\begin{aligned} \frac{W_n(\eta'_x \sigma)}{W_n(\sigma)} &= \exp(2a(n)k(\eta') - 2b(n)(\gamma(\eta'_x \sigma) - \gamma(\sigma))) \\ &\leq \exp((2a(n) + 2\mathcal{V}b(n))k(\eta')) . \end{aligned}$$

Now, the hypothesis $a(n) + \mathcal{V}b(n) \leq 0$ implies the ratio $W_n(\eta'_x \sigma)$ divided by $W_n(\sigma)$ is smaller than 1. Thanks to (8), the conditional probability $\mu_{a,b}(I_x^\eta = 1 | \sigma)$ is larger than $|\mathcal{C}_r|^{-1} W_n(\eta)$; $c_r = |\mathcal{C}_r|^{-1} = 2^{-\beta_r}$ is suitable.

Since the lower bound of (6) is uniform on the configuration $\sigma \in \{-, +\}^{\delta B(x, r)}$, the same inequality holds for

$$\mu_{a,b}(I_x^\eta = 1) = \mathbb{E}_{a,b}[\mu_{a,b}(I_x^\eta = 1 | \mathcal{F}(\delta B(x, r)))] .$$

Finally, (7) has been proved in Proposition 3.2 of [8] with $C_r = 2^{\mathcal{V}\beta_r}$. $\square$

The lower bound given by (6) has the advantage of being uniform on the configuration $\sigma \in \{-, +\}^{\delta B(x,r)}$. This is the reason why this inequality will be often used in the proofs. In addition, there is no uniform upper bound for the conditional probability $\mu_{a,b}(I_x^\eta = 1|\sigma)$. In order to make up for this gap, we will have recourse to the FKG-inequality.

There is a natural partial ordering on the configuration set $\mathcal{X}_n = \{-, +\}^{V_n}$ defined by $\sigma \leq \sigma'$ if $\sigma(x) \leq \sigma'(x)$ for all vertices $x \in V_n$. A real function defined on $\mathcal{X}_n$ is *increasing* if $f(\sigma) \leq f(\sigma')$ whenever $\sigma \leq \sigma'$. An event $A \in \mathcal{F}(\mathcal{X}_n)$ is also said *increasing* whenever its indicator function $f = \mathbb{1}_A$ is increasing. Conversely, a *decreasing* event is an event whose complementary set (in $\mathcal{F}(\mathcal{X}_n)$) is increasing.

Let us focus on an example of increasing event which will be central in our study. For $0 \leq W \leq 1$, we denote by $\mathcal{C}_r(W)$ the set of local configurations with radius r whose weight is smaller than W :

$$\mathcal{C}_r(W) = \{\eta \in \mathcal{C}_r, W_n(\eta) \leq W\} .$$

Thus, let us consider two local configurations η, η' such that the set of positive vertices of η' contains that of η . The perimeter $\gamma(\eta')$ is not necessary larger than $\gamma(\eta)$: roughly, $V_+(\eta)$ may have holes. However, the inequality

$$\gamma(\eta') \geq \gamma(\eta) - (k(\eta') - k(\eta))\mathcal{V}$$

holds. Hence, the ratio

$$\begin{aligned} \frac{W_n(\eta')}{W_n(\eta)} &= \exp(2a(n)(k(\eta') - k(\eta)) - 2b(n)(\gamma(\eta') - \gamma(\eta))) \\ &\leq \exp((2a(n) + 2\mathcal{V}b(n))(k(\eta') - k(\eta))) \end{aligned}$$

is smaller than 1 whenever $a(n) + \mathcal{V}b(n)$ is negative. As a consequence, under this hypothesis, the set $\mathcal{C}_r(W)$ allows to build some increasing variables; for instance, the random indicator

$$\sum_{\eta \in \mathcal{C}_r(W)} I_x^\eta .$$

For a positive value of the pair potential b , the Gibbs measure $\mu_{a,b}$ defined by (2) satisfies the FKG inequality, i.e.

$$\mu_{a,b}(A \cap A') \geq \mu_{a,b}(A)\mu_{a,b}(A') , \quad (9)$$

for all increasing events A and A' : see for instance Section 3 of [15]. In this case, the Ising model is said *ferromagnetic*. Note that inequality (9) can be immediately extended to decreasing events (see [18] p.). Moreover, the union and the intersection of increasing events are still increasing. The same is true for decreasing events. Hence, the FKG inequality applies to any family made up of a finite number of decreasing events $A_1, \dots, A_m$:

$$\mu_{a,b}(A_1 \cap \dots \cap A_m) \geq \mu_{a,b}(A_1) \dots \mu_{a,b}(A_m) . \quad (10)$$

4 Exponential bounds for the probability of nonexistence

Let $x \in V_n$ be a vertex, $R \geq r \geq 1$ be two integers and W be a positive real. Let us denote by $\mathcal{A}(x, R, W)$ the following event:

$$\exists y \in B(x, R - r), \exists \eta \in \mathcal{C}_r(W), I_y^\eta = 1 .$$

The event $\mathcal{A}(x, R, W)$ means at least one copy of a local configuration (with radius r) whose weight is smaller than W can be found somewhere in the large ball $B(x, R)$. Theorem 4.1 gives exponential bounds for the probability of the complementary event $\mathcal{A}(x, R(n), W_n)^c$.

Theorem 4.1 *Assume that the magnetic field $a(n)$ is negative, the pair potential $b(n)$ is nonnegative and they satisfy $a(n) + 2\mathcal{V}b(n) \leq 0$. Let $(W_n)_{n \in \mathbb{N}}$ be a sequence of positive reals satisfying the following property: there exist an integer N and $0 < \epsilon < 1$ such that*

$$\forall n \geq N, \min_{\eta \in \mathcal{C}_r} W_n(\eta) \leq W_n \leq \epsilon c_r C_r^{-1} .$$

Let $K = (1 - \epsilon)^{-1} c_r^{-1} C_r$ and $K' = \tau c_r$. Then, for all $n \geq N$ and for all vertex x ,

$$e^{-KR(n)^d W_n} \leq 1 - \mu_{a,b}(\mathcal{A}(x, R(n), W_n)) \leq e^{-K'R(n)^d W_n} . \quad (11)$$

Remark the constants K and K' only depend on r, ϵ and parameters q, ρ and d (the constant τ will be introduced in the proof).

The fact that, for n large enough, W_n is assumed larger than the smallest weight $W_n(\eta)$, $\eta \in \mathcal{C}_r$, only serves to ensure that the set $\mathcal{C}_r(W_n)$ (and the event $\mathcal{A}(x, R(n), W_n)$ too) is non empty. Moreover, let us note hypothesis $W_n \leq \epsilon c_r C_r^{-1}$ is only used in the proof of the lower bound of (11).

The inequalities of (11) give a limit for the probability of $\mathcal{A}(x, R(n), W_n)$ depending on whether the product $R(n)^d W_n$ tends to 0 or $+\infty$:

$$\text{if } \lim_{n \rightarrow +\infty} R(n)^d W_n = 0 \text{ then } \lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{A}(x, R(n), W_n)) = 0 ; \quad (12)$$

$$\text{if } \lim_{n \rightarrow +\infty} R(n)^d W_n = +\infty \text{ then } \lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{A}(x, R(n), W_n)) = 1 . \quad (13)$$

Roughly speaking, if the radius $R(n)$ is small compared to $W_n^{-1/d}$ then asymptotically, with probability tending to 1, there is no local configuration whose weight is smaller than W_n . Conversely, if $R(n)$ is large compared to $W_n^{-1/d}$ then at least one copy of an element of $\mathcal{C}_r(W_n)$ can be found somewhere in $B(x, R(n))$, with probability tending to 1. Relation (12) and (13) complete Proposition 4.2 of [11].

Finally, Theorem 4.1 implies the quantity

$$\frac{1}{R(n)^d W_n} \ln (\mu_{a,b}(\mathcal{A}(x, R(n), W_n)^c)) \quad (14)$$

belongs to the interval $[-K, -K']$. A natural question is to wonder whether this ratio admits a limit as n goes to infinity. It seems to be difficult to answer to this question in general. However, in the following particular case, the answer is positive and the corresponding limit is known; see [9] or [10] for more details. Assume the weight $W_n = W_n(\eta)$ is the one of a local configuration η , the pair potential $b(n) = b$ is a positive real number and the magnetic field $a(n)$ tends to $-\infty$. Under these conditions, the limit of the probability for a given local configuration to occur on the ball $B(x, R(n))$ essentially depends on its number of positive vertices. Moreover, assume the radius $R(n)$ is such that the product $R(n)^d W_n = R(n)^d W_n(\eta)$ is constant. Then, the quantity (14) converges, as $n \rightarrow +\infty$, to

$$- \sum_{\substack{\eta' \in \mathcal{C}_r \\ k(\eta') = k(\eta)}} e^{-2b(\gamma(\eta') - \gamma(\eta))} .$$

This section ends with the proof of Theorem 4.1.

Proof: Throughout this proof, potentials $a(n)$ and $b(n)$, and the sequence $(W_n)_{n \in \mathbb{N}}$ satisfy the hypotheses of Theorem 4.1.

The two following remarks will lighten notations and formulas of the proof. Thanks to the invariance translation of the graph G_n it suffices to prove Theorem 4.1 for $x = 0$. Hence, we merely denote by $\mathcal{A}(R(n), W_n)$ the event $\mathcal{A}(0, R(n), W_n)$. Furthermore, $\mathcal{A}(R(n), W_n)$ and $\mathcal{A}(R(n), W'_n)$ are equal for

$$W'_n = \max_{\eta \in \mathcal{C}_r(W_n)} W_n(\eta) .$$

As a consequence and without loss of generality, we can assume that, for each n , the weight W_n belongs to $\{W_n(\eta), \eta \in \mathcal{C}_r\}$.

The proof of the lower bound of (11) requires the ferromagnetic character of the Ising model, i.e. the positivity of the pair potential $b(n)$. Since $a(n) + \mathcal{V}b(n)$ is negative, the event

$$\bigcap_{\eta \in \mathcal{C}_r(W_n)} \{I_y^\eta = 0\}$$

is decreasing, whatever the vertex y . So the FKG inequality implies:

$$\mu_{a,b}(\mathcal{A}(R(n), W_n)^c) \geq \prod_{y \in B(0, R(n) - r)} \mu_{a,b} \left(\bigcap_{\eta \in \mathcal{C}_r(W_n)} \{I_y^\eta = 0\} \right) . \quad (15)$$

Now, it suffices to control each term of the above product. This is the role of Lemma 3.2. Let us pick $y \in B(0, R(n) - r)$. We get:

$$\begin{aligned} \mu_{a,b} \left(\bigcap_{\eta \in \mathcal{C}_r(W_n)} \{I_y^\eta = 0\} \right) &= 1 - \sum_{\eta \in \mathcal{C}_r(W_n)} \mu_{a,b}(I_y^\eta = 1) \\ &\geq 1 - \sum_{\eta \in \mathcal{C}_r(W_n)} C_r W_n(\eta) \\ &\geq 1 - c_r^{-1} C_r W_n , \end{aligned}$$

since $|\mathcal{C}_r(W_n)| \leq |\mathcal{C}_r| = c_r^{-1}$. Let N be the integer introduced in the statement of Theorem 4.1 and let $n \geq N$. Therefore, from (15) and the inequality $\ln(1 - X) \geq -(1 - \epsilon)^{-1}X$ valid for $0 \leq X \leq \epsilon$, it follows that:

$$\begin{aligned} \mu_{a,b}(\mathcal{A}(R(n), W_n)^c) &\geq (1 - c_r^{-1}C_r W_n)^{R(n)^d} \\ &\geq \exp(-(1 - \epsilon)^{-1}c_r^{-1}C_r R(n)^d W_n) . \end{aligned}$$

So, the lower bound of (11) is proved with $K = (1 - \epsilon)^{-1}c_r^{-1}C_r$.

In order to prove the upper bound of (11), let us denote by T_n the subset of V_n defined by:

$$T_n = \left\{ i(\rho(2r+1)+1), |i| = 0, \dots, \left\lfloor \frac{\rho R(n)}{\rho(2r+1)+1} \right\rfloor - 1 \right\}^d \cap B(0, R(n) - r) .$$

Its cardinality, denoted by τ_n , satisfies $\tau_n \geq \tau R(n)^d$, for a positive constant $\tau > 0$ not depending on the size n . Let $\mathcal{T}_n$ be the union of the balls of radius r centered at the elements of T_n (see Figure 2)

$$\mathcal{T}_n = \bigcup_{y \in T_n} B(y, r)$$

and denote by X_n the random variable which counts the number of copies of elements of $\mathcal{C}_r(W_n)$ occurring on balls $B(y, r)$, $y \in T_n$,

$$X_n = \sum_{y \in T_n} \sum_{\eta \in \mathcal{C}_r(W_n)} I_y^\eta .$$




Figure 2: The colored vertices represent the set $\mathcal{T}_n$ in dimension $d = 2$, with $r = 2$, $\rho = 2$ and relative to the L_1 norm. The black vertices are elements of T_n

Since the balls $B(y, r)$, $y \in T_n$, are included in $B(0, R(n))$, the event $X_n > 0$ implies $\mathcal{A}(R(n), W_n)$. So, the upper bound of (11) follows from the next statement. Let $K' = \tau c_r$. Then, for all n ,

$$\mu_{a,b}(X_n = 0) \leq e^{-K'R(n)^d W_n} . \quad (16)$$

Let us prove this inequality. Remark that any two balls $B(y, r)$ and $B(y', r)$ where y and y' are distinct vertices of T_n are $\mathcal{V}$ -disjoint. Hence, Property 3.1 produces the following identities:

$$\begin{aligned}
\mu_{a,b}(X_n = 0) &= \mathbb{E}_{a,b}[\mu_{a,b}(X_n = 0 \mid \mathcal{F}(\delta\mathcal{T}_n))] \\
&= \mathbb{E}_{a,b}\left[\mu_{a,b}\left(\bigcap_{y \in T_n} \bigcap_{\eta \in \mathcal{C}_r(W_n)} I_y^\eta = 0 \mid \mathcal{F}(\delta\mathcal{T}_n)\right)\right] \\
&= \mathbb{E}_{a,b}\left[\prod_{y \in T_n} \mu_{a,b}\left(\bigcap_{\eta \in \mathcal{C}_r(W_n)} I_y^\eta = 0 \mid \mathcal{F}(\delta\mathcal{T}_n)\right)\right] \\
&= \mathbb{E}_{a,b}\left[\prod_{y \in T_n} \mu_{a,b}\left(\bigcap_{\eta \in \mathcal{C}_r(W_n)} I_y^\eta = 0 \mid \mathcal{F}(\delta B(y, r))\right)\right]. \tag{17}
\end{aligned}$$

Now, let $y \in T_n$ be a vertex and $\sigma \in \{-, +\}^{\delta B(y, r)}$ a configuration. We can write:

$$\begin{aligned}
\mu_{a,b}\left(\bigcap_{\eta \in \mathcal{C}_r(W_n)} I_y^\eta = 0 \mid \sigma\right) &= 1 - \mu_{a,b}\left(\bigcup_{\eta \in \mathcal{C}_r(W_n)} I_y^\eta = 1 \mid \sigma\right) \\
&\leq 1 - \mu_{a,b}(I_y^\eta = 1 \mid \sigma), \tag{18}
\end{aligned}$$

where η is an element of $\mathcal{C}_r(W_n)$ satisfying $W_n(\eta) = W_n$. Lemma 3.2 gives a bound for (18) which does not depend on the configuration $\sigma \in \{-, +\}^{\delta B(y, r)}$ nor on the vertex y . Hence, it follows from (17):

$$\mu_{a,b}(X_n = 0) \leq (1 - c_r W_n)^{\tau_n}$$

(recall that $1 - c_r W_n$ is positive). Finally, using $\tau_n \geq \tau R(n)^d$ and the classical inequality $\ln(1 + X) \leq X$ valid for $X > -1$, a bound for the probability of X_n to be null is obtained:

$$\mu_{a,b}(X_n = 0) \leq \exp(-\tau R(n)^d c_r W_n).$$

The inequality (16) is proved with $K' = \tau c_r$. □

5 Distance between local configurations

5.1 Motivation

Throughout Section 5, η represents a local configuration with radius r having at least one positive vertex, i.e. different from the null local configuration η^0 . The goal of this section

is to describe the Ising model under the hypothesis (H) , bearing on the potentials $a(n)$ and $b(n)$, and defined by:

$$(H) : \begin{cases} n^d W_n(\eta) = \lambda, & \text{for } \lambda > 0 \\ a(n) + 2\mathcal{V}b(n) \leq 0 \\ a(n) + \mathcal{V}b(n) \rightarrow -\infty & \text{as } n \rightarrow +\infty \end{cases}$$

Let us start by giving the reason for (H) . Let $X_n(\eta)$ be the random variable which counts the number of copies of η in the whole graph G_n :

$$X_n(\eta) = \sum_{x \in V_n} I_x^\eta .$$

In [8], various results have been stated about the variable $X_n(\eta)$. If the product $n^d W_n(\eta)$ tends to 0 (resp. $+\infty$) then the probability $\mu_{a,b}(X_n(\eta) > 0)$ tends to 0 (resp. 1). In other words, if $W_n(\eta)$ is small compared to n^{-d} , then asymptotically, there is no occurrence of η in G_n . If $W_n(\eta)$ is large compared to n^{-d} , then at least one occurrence of η can be found in the graph, with probability tending to 1. Moreover, provided the hypothesis (H) is satisfied (in particular $W_n(\eta)$ is of order n^{-d}), the distribution of $X_n(\eta)$ converges weakly to the Poisson distribution with parameter λ .

In particular, the number of copies of η occurring in the graph is finite with probability tending to 1. Let η_x be one of them (that occurring on the ball centered at x). Lemma 5.1 says vertices around η_x are negative with probability tending to 1. For that purpose, let us introduce the *ring* $\mathcal{R}(x, r, R)$ defined as the following set of vertices

$$\mathcal{R}(x, r, R) = \{y \in V_n, r < \text{dist}(x, y) \leq R\} .$$

Lemma 5.1 *Let η be a local configuration with radius r and let $x \in V_n$. Let us denote by $\mathcal{A}_r(x, R, +)$ the (interpreted) event “there exists at least one positive vertex in $\mathcal{R}(x, r, R)$ ”. If the potentials satisfy $a(n) + 2\mathcal{V}b(n) \leq 0$ and $a(n) \rightarrow -\infty$. Then,*

$$\lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{A}_r(x, R, +) \mid I_x^\eta = 1) = 0 .$$

Proof: Let us introduce the set $\mathcal{C}_R(\eta, +)$ formed from local configurations with radius R whose restriction to the ball $B(0, r)$ is equal to η and having at least one positive vertex in the ring $\mathcal{R}(0, r, R)$:

$$\mathcal{C}_R(\eta, +) = \{\zeta \in \mathcal{C}_R, \forall y \in B(0, r), \zeta(y) = \eta(y) \text{ and } k(\zeta) > k(\eta)\} .$$

The intersection of the events $\mathcal{A}_r(x, R, +)$ and $I_x^\eta = 1$ forces the ball $B(x, R)$ to contain at least $k(\eta) + 1$ positive vertices. Precisely,

$$\mathcal{A}_r(x, R, +) \cap \{I_x^\eta = 1\} \subset \bigcup_{\zeta \in \mathcal{C}_R(\eta, +)} \{I_x^\zeta = 1\} .$$

Then, Lemma 3.2 allows us to bound the probability of that intersection:

$$\begin{aligned} \mu_{a,b}(\mathcal{A}_r(x, R, +) \cap \{I_x^\eta = 1\}) &\leq \sum_{\zeta \in \mathcal{C}_R(\eta, +)} \mu_{a,b}(I_x^\zeta = 1) \\ &\leq C_R \sum_{\zeta \in \mathcal{C}_R(\eta, +)} W_n(\zeta) . \end{aligned} \quad (19)$$

Let ζ be an element of $\mathcal{C}_R(\eta, +)$. By definition, $k(\zeta) \geq k(\eta) + 1$. Moreover, thanks to the convexity of balls, the perimeter of ζ is necessary as large as that of η . In other words, $W_n(\zeta) \leq e^{2a(n)} W_n(\eta)$. Finally, we deduce from (19) that the conditional probability satisfies

$$\mu_{a,b}(\mathcal{A}_r(x, R, +) \mid I_x^\eta = 1) \leq c_r^{-1} C_R |\mathcal{C}_R(\eta, +)| e^{2a(n)}$$

and tends to 0 as n tends to infinity. $\square$

This result constitutes the starting point of our study since from now on, a natural question concerns the distance from η_x to the closest positive vertices. Theorems 5.3 and 5.4 will answer to this question and, more generally, will describe the distance between local configurations under (H) .

5.2 The main results

The main results of Section 5, Theorems 5.3 and 5.4, concern the two following events.

Definition 5.2 *Let $r \geq 1$, $R \geq 3r + 1$, $0 < W < 1$ and $x \in V_n$. Let us denote by $\mathcal{A}_r(x, R, W)$ the event*

$$\exists y \in \mathcal{R}(x, 2r, R - r), \exists \zeta \in \mathcal{C}_r(W) \text{ such that } I_y^\zeta = 1$$

and, for all integer $\ell > 2r$, by $\mathcal{H}_r(x, R, \ell, W)$ the event

$$\begin{aligned} \exists y, y' \in \mathcal{R}(x, 2r, R - r), \exists \zeta, \zeta' \in \mathcal{D}_r(W) \text{ such that} \\ 2r < \text{dist}(y, y') \leq \ell \text{ and } I_y^\zeta = I_{y'}^{\zeta'} = 1 , \end{aligned}$$

where $\mathcal{D}_r(W)$ is the set of local configurations with radius r whose weight is equal to W .

The event $\mathcal{A}_r(x, R, W)$ means a local configuration with radius r whose weight is smaller than W can be found somewhere in the ring $\mathcal{R}(x, r, R)$. It generalizes the notation $\mathcal{A}_r(x, R, +)$ of Lemma 5.1. Indeed, a single positive vertex can be viewed as a local configuration with radius r and weight $W = e^{2a(n) - 2Vb(n)}$.

The event $\mathcal{H}_r(x, R, \ell, W)$ corresponds to the occurrence in $\mathcal{R}(x, r, R)$ of two local configurations of weight W at distance from each other smaller than ℓ . The precaution $\text{dist}(y, y') > 2r$ ensures that the occurrences of ζ and ζ' do not use the same positive vertices.

Theorem 4.1 says whatever the choice of radii $R \geq r$, the probability for the null local

configuration η^0 (i.e. the element of $\mathcal{C}_r$ having only negative vertices) to occur in the ball $B(x, R)$ is positive:

$$\mu_{a,b}(\mathcal{A}(x, R, 1)) \geq 1 - e^{-K'R^d} .$$

This being said, we focus our attention on local configurations having at least one positive vertex. Now, the hypothesis (H) forces the weight of such an element to tend to 0 as n tends to infinity. Hence, the study of the conditional probability $\mu_{a,b}(\mathcal{A}_r(x, R, W) | I_x^\eta = 1)$ becomes relevant when the weight $W = W_n$ is allowed to depend on the size n and to tend to 0. Moreover, in order to avoid trivial situations, it is reasonable to assume that $\mathcal{C}_r(W_n)$ is a nonempty set. Now, remark the events $\mathcal{A}_r(x, R, W_n)$ and $\mathcal{A}_r(x, R, W'_n)$ are equal for

$$W'_n = \max_{\zeta \in \mathcal{C}_r(W_n)} W_n(\zeta) .$$

As a consequence and without loss of generality, we can assume that, for each n , the weight W_n belongs to $\{W_n(\zeta), \zeta \in \mathcal{C}_r\}$. From then on, Lemma 5.1 implies

$$\mu_{a,b}(\mathcal{A}_r(x, R, W_n) | I_x^\eta = 1) \rightarrow 0 ,$$

for any fixed radius R . So, in order to obtain a positive limit for the previous quantity, it is needed to take a radius $R = R(n)$ which tends to $+\infty$. All these remarks are still true for $\mathcal{H}_r(x, R, \ell, W)$.

To sum up, the sequences $(R(n))_{n \in \mathbb{N}}$ and $(W_n)_{n \in \mathbb{N}}$ will be assumed to satisfy the hypothesis (H'):

$$(H') : \begin{cases} R(n) \rightarrow +\infty \\ W_n \rightarrow 0 \\ \forall n, \exists \zeta_n \in \mathcal{C}_r, W_n = W_n(\zeta_n) \end{cases}$$

Imagine a copy of η occurs on $B(x, r)$. Theorem 5.3 says that the distance to η_x from the closest positive vertices is of order

$$(e^{2a(n)-2\mathcal{V}b(n)})^{-1/d}$$

and, more generally, the distance to η_x from the closest local configurations of weight W_n is of order $W_n^{-1/d}$.

Theorem 5.3 *Let us consider a local configuration $\eta \in \mathcal{C}_r$ and potentials $a(n) < 0$ and $b(n) \geq 0$ satisfying (H). Let $(R(n))_{n \in \mathbb{N}}$ and $(W_n)_{n \in \mathbb{N}}$ be two sequences satisfying (H'). Let x be a vertex. Then,*

$$\lim_{n \rightarrow +\infty} R(n)W_n^{1/d} = 0 \implies \lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{A}_r(x, R(n), W_n) | I_x^\eta = 1) = 0 \quad (20)$$

and

$$\lim_{n \rightarrow +\infty} R(n)W_n^{1/d} = +\infty \implies \lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{A}_r(x, R(n), W_n) | I_x^\eta = 1) = 1 . \quad (21)$$

First, let us precise that the second part of Theorem 5.3 concerns only the case where the speed of convergence to 0 of the weight W_n is slower than that of $W_n(\eta)$. Indeed, the radius $R(n)$ can not exceed the size n of the graph G_n and the product $n^d W_n(\eta)$ is constant.

Let us describe Theorem 5.3 when potentials $a(n)$ and $b(n)$ satisfy the following hypotheses:

$$a(n) \rightarrow -\infty, b(n) \rightarrow +\infty \text{ and } \frac{b(n)}{a(n)} \rightarrow 0 .$$

Assume a copy of η occurs on $B(x, r)$ and consider two local configurations η_1 and η_2 (whose weights are larger than that of η). Then, this is first the number of positive vertices thus the perimeter which allow to know what are the closest to η_x , among the copies of η_1 and η_2 . Indeed, if $k(\eta_1) < k(\eta_2) (\leq k(\eta))$ then the distance from the closest copies of η_1 to η_x is smaller than the distance from the closest copies of η_2 to η_x , whatever their perimeters. Furthermore, if η_1 and η_2 have the same number of positive vertices then that having the smallest perimeter will have its closest copies (to η_x) closer to η_x than the closest copies (to η_x) of the other one. Figure 3 proposes a panorama of this situation.




Figure 3: Let $\rho = 1$, $q = \infty$ and $d = 2$; each vertex has $\mathcal{V} = 8$ neighbors. Assume potentials satisfy the hypothesis (H), $a(n) \rightarrow -\infty$ and the ratio $b(n)/a(n) \rightarrow 0$. A copy of η occurs on the ball $B(x, r)$. First, the closest local configurations to η_x are represented according to their number of positive vertices. Thus, the situation of those having $k = 3$ positive vertices is precised. The closest ones to η_x are those having a perimeter $\gamma = 18$ and the farthest ones are those having isolated positive vertices, i.e $\gamma = 24$.

Now, let us consider two copies of a given local configuration ζ which are at distance to η_x of order $W_n(\zeta)^{-1/d}$. Then, the distance between these two copies necessarily tends to infinity. Otherwise, they would form together a “super” local configuration whose weight would be smaller than $W_n(\zeta)$ and which would be at distance of order $W_n(\zeta)^{-1/d}$ to η_x . This would contradict Theorem 5.3. Theorem 5.4 precises this situation; it says the distance between these two copies of ζ tends to infinity as $W_n(\zeta)^{-1/d}$.

Before stating Theorem 5.4, let us introduce the *index* of W_n . The hypothesis (H') ensures the set $\mathcal{D}_r(W_n)$ is nonempty. However, it is not necessary reduced to only one element;

it may even contain some local configurations not having the same number of positive vertices. Hence, let us denote by k_n and call the index of W_n this maximal number:

$$k_n = \max_{\zeta \in \mathcal{D}_r(W_n)} k(\zeta) .$$

Theorem 5.4 *Let us consider a local configuration $\eta \in \mathcal{C}_r$ and potentials $a(n) < 0$ and $b(n) \geq 0$ satisfying (H). Let $(R(n))_{n \in \mathbb{N}}$ and $(W_n)_{n \in \mathbb{N}}$ be two sequences satisfying (H'). Let $(\ell(n))_{n \in \mathbb{N}}$ be a sequence of integers and x be a vertex. Then,*

$$\begin{aligned} \lim_{n \rightarrow +\infty} R(n) W_n^{2/d} \max \left\{ \ell(n), e^{\frac{2}{d} \mathcal{V} b(n) k_n} \right\} &= 0 \\ \implies \lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{H}_r(x, R(n), \ell(n), W_n) \mid I_x^\eta = 1) &= 0 \end{aligned} \quad (22)$$

and

$$\lim_{n \rightarrow +\infty} R(n) \ell(n) W_n^{2/d} = +\infty \implies \lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{H}_r(x, R(n), \ell(n), W_n) \mid I_x^\eta = 1) = 1 . \quad (23)$$

Assume a copy of η occurs on $B(x, r)$. We know (Theorem 5.3) the distance from η_x to the closest (copies of) local configurations of weight W_n is of order $W_n^{-1/d}$. Let us suppose the radius $R(n)$ is of order $W_n^{-1/d}$ (i.e. $R_n^d W_n$ tends to a positive constant) and let ζ_y be such a copy (i.e. $\zeta \in \mathcal{D}_r(W_n)$ occurs on $B(y, r)$ and $\text{dist}(x, y)$ is of order $R(n)$). If $\ell(n)$ is large compared to $W_n^{-1/d}$, then the product $R(n) \ell(n) W_n^{2/d}$ tends to infinity and (23) says a copy of a local configuration of weight W_n can be found at distance from ζ_y smaller than $\ell(n)$. Conversely, if $\ell(n)$ is small compared to $W_n^{-1/d}$ then

$$R(n) W_n^{2/d} \max \left\{ \ell(n), e^{\frac{2}{d} \mathcal{V} b(n) k_n} \right\} \rightarrow 0 .$$

Indeed,

$$\begin{aligned} W_n e^{2 \mathcal{V} b(n) k_n} &= W_n e^{-2a(n) k_n} e^{2k_n(a(n) + \mathcal{V} b(n))} \\ &\leq e^{a(n) + \mathcal{V} b(n)} \end{aligned}$$

which tends to 0 as $n \rightarrow +\infty$ thanks to the hypothesis (H). From then on, (22) says there is no copy of local configurations of weight W_n at distance from ζ_y smaller than $\ell(n)$. Figure 4 represents various local configurations and the distances between each other.

Besides, if the product $R(n) W_n^{2/d}$ tends to infinity then (23) implies that, at distance from η_x of order $R(n)$, one can find two copies of local configurations of weight W_n so close to each other that we wish. This is not surprising: Theorem 5.3 claims there are local configurations of weight W_n^2 in the ring $\mathcal{R}(x, r, R(n))$ provided $R(n) W_n^{2/d} \rightarrow +\infty$.

Furthermore, let us underline that Theorems 5.3 and 5.4 would remain unchanged if the radius of local configurations occurring in $\mathcal{R}(x, r, R(n))$ might be different from that of η .

Let us end this section by the following remark. Assume that the local configuration η occurs on $B(x, r)$. As long as the ratio $R(n)/n$ tends to 0, Theorem 5.3 says there is




Figure 4: Let η be the local configuration having $k(\eta) = 5$ positive vertices represented above. Assume potentials $a(n)$ and $b(n)$ satisfy (H) . The distance from a given copy of η occurring in the graph, say η_x , to its closest positive vertices is of order $W_n(+)^{-1/d}$ where $W_n(+)$ denotes the weight of a single positive vertex. The distance between such positive vertices is of order $W_n(+)^{-1/d}$ too. Let ζ be the local configuration having $k(\zeta) = 3$ positive vertices represented above. The distance from the closest copies of ζ to η_x is of order $W_n(\zeta)^{-1/d}$ and the distance between such copies is $W_n(\zeta)^{-1/d}$ too. Finally, the other copies of η occurring in G_n are at distance to η_x of order n , since $n^d W_n(\eta) = \lambda$.

no other copy of η in the ring $\mathcal{R}(x, r, R(n))$. Now, n represents the size of the graph G_n . So, if other copies (than η_x) of η occur in the graph, they should be at distance of order n from η_x . This is the meaning of Proposition 5.5. Recall that $X_n(\eta)$ represents the number of copies of η in G_n . We denote by $\mathcal{E}_n(\eta)$ the event

$$\mathcal{E}_n(\eta) : \exists C > 0, (\forall x \neq y, I_x^\eta = I_y^\eta = 1) \implies \text{dist}(x, y) \geq Cn .$$

Proposition 5.5 *Let us consider a local configuration $\eta \in \mathcal{C}_r$ and potentials $a(n) < 0$ and $b(n) \geq 0$ satisfying (H) . Then,*

$$\lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{E}_n(\eta) \mid X_n(\eta) \geq 2) = 1 .$$

Let us recall that the random variable $X_n(\eta)$ converges weakly to a Poisson distribution when the hypothesis (H) is satisfied (see [8]). In particular, the fact that no more than one copy of η (i.e 0 or 1) occurs in the graph has a positive asymptotic probability. So, conditioning by $X_n(\eta) \geq 2$, we avoid this uninteresting case.

5.3 Proofs of Theorems 5.3 and 5.4 and Proposition 5.5

The intuition behind Theorem 5.3 is the following. On the one hand, the Markovian character of the Gibbs measure $\mu_{a,b}$ implies that the events $\mathcal{A}_r(x, R(n), W_n)$ and $\{I_x^\eta = 1\}$ can be considered as asymptotically independent. So, as n goes to infinity, $\mu_{a,b}(\mathcal{A}_r(x, R(n), W_n) | I_x^\eta = 1)$ and $\mu_{a,b}(\mathcal{A}_r(x, R(n), W_n))$ evolve in the same way. On the other hand, the events $\mathcal{A}_r(x, R(n), W_n)$ and $\mathcal{A}(x, R(n), W_n)$ (see Section 4) are the same but for a finite number of vertices, those belonging to $B(x, r)$. Their probabilities have the same limit. In conclusion, Theorem 4.1 implies the conditional probability $\mu_{a,b}(\mathcal{A}_r(x, R(n), W_n) | I_x^\eta = 1)$ should tend to 0 or 1 according to the quantity $R(n)^d W_n$. The same remarks hold for Theorem 5.4.

Thanks to the invariance translation of the graph G_n , we only prove Theorem 5.3 and 5.4 for $x = 0$ and we will denote respectively by $\mathcal{A}_r(R(n), W_n)$ and $\mathcal{H}_r(R(n), \ell(n), W_n)$ the events $\mathcal{A}_r(0, R(n), W_n)$ and $\mathcal{H}_r(0, R(n), \ell(n), W_n)$.

Proof (of Theorem 5.3): Recall the definition of $\mathcal{A}_r(R(n), W_n)$:

$$\exists y \in \mathcal{R}(0, 2r, R(n) - r), \exists \zeta \in \mathcal{C}_r(W_n), I_y^\zeta = 1 .$$

Let us start with the proof of (20). The first step is to move the occurrence of ζ (the local configuration fulfilling $\mathcal{A}_r(R(n), W_n)$) away from the ball $B(0, r)$ on which occurs η . Let y be a vertex belonging to the ring $\mathcal{R}(0, 2r, R - r)$. Either $\text{dist}(y, 0) \leq 2r + 1$ and the ball $B(y, r)$ is included in $\mathcal{R}(0, r, 3r + 1)$. Either $\text{dist}(y, 0) > 2r + 1$ and $B(y, r)$ is included in $\mathcal{R}(0, r + 1, R(n))$. In other words,

$$\mathcal{A}_r(R(n), W_n) \subset \mathcal{A}_r(3r + 1, W_n) \cup \mathcal{A}_{r+1}(R(n), W_n) . \quad (24)$$

The hypothesis (H') forces the weight W_n to be smaller than $e^{2a(n) - 2\mathcal{V}b(n)}$. Hence, the event $\mathcal{A}_r(3r + 1, W_n)$ is included in $\mathcal{A}_r(0, 3r + 1, +)$ and Lemma 5.1 implies that the conditional probability

$$\mu_{a,b}(\mathcal{A}_r(3r + 1, W_n) | I_0^\eta = 1)$$

tends to 0. From (24), it remains to prove the same limit holds for

$$\mu_{a,b}(\mathcal{A}_{r+1}(R(n), W_n) | I_0^\eta = 1) .$$

The strategy consists in introducing increasing events in order to use the FKG inequality. Since $a(n) + \mathcal{V}b(n)$ is negative, the following events are both increasing:

$$\bigcup_{\zeta \in \mathcal{C}_r(W_n)} \{I_y^\zeta = 1\} \quad \text{and} \quad \bigcup_{\eta' \in \mathcal{C}_r(W_n(\eta))} \{I_0^{\eta'} = 1\} .$$

So does their intersection

$$S_y = \left(\bigcup_{\zeta \in \mathcal{C}_r(W_n)} \{I_y^\zeta = 1\} \right) \cap \left(\bigcup_{\eta' \in \mathcal{C}_r(W_n(\eta))} \{I_0^{\eta'} = 1\} \right) .$$

As a consequence, we get from the FKG inequality:

$$1 - \mu_{a,b}(\mathcal{A}_{r+1}(R(n), W_n) \cap \{I_0^\eta = 1\}) \geq \prod_{y \in \mathcal{R}(0, 2r+1, R(n)-r)} \mu_{a,b}(S_y^c) . \quad (25)$$

Let $y \in \mathcal{R}(0, 2r+1, R(n)-r)$. Let us denote by V_y the union of the two balls $B(y, r)$ and $B(0, r)$, and by $\mathcal{C}_y$ the set of configurations defined below:

$$\mathcal{C}_y = \{\omega \in \{-, +\}^{V_y}, W_n(\omega_{B(0,r)}) \leq W_n(\eta) \text{ and } W_n(\omega_{B(y,r)}) \leq W_n\} .$$

Then, the probability $\mu_{a,b}(S_y^c)$ becomes

$$\mu_{a,b}(S_y^c) = 1 - \sum_{\omega \in \mathcal{C}_y} \mu_{a,b}(I_{V_y}^\omega = 1) ,$$

where $I_{V_y}^\omega$ is the random indicator defined on $\mathcal{X}_n$ as follows: $I_{V_y}^\omega(\sigma)$ is 1 if the restriction of the configuration $\sigma \in \mathcal{X}_n$ to the set V_y is ω and 0 otherwise. Let ω be an element of $\mathcal{C}_y$. Actually, the inequality (7) of Lemma 3.2 can be extended to any subset of vertices V (see Proposition 3.2 of [8]): whenever $a(n) + 2Vb(n)$ is negative, there exists a constant $C_V > 0$ such that for all $\omega \in \{-, +\}^V$,

$$\mu_{a,b}(I_V^\omega = 1) \leq C_V W_n(\omega) . \quad (26)$$

Moreover, C_V depends on the set V only through its cardinality. Applying (26) to V_y , we obtain:

$$\mu_{a,b}(S_y^c) \geq 1 - \sum_{\omega \in \mathcal{C}_y} C_{2,r} W_n(\omega) ,$$

where the constant $C_{2,r}$ only depends on r . Now, $\text{dist}(0, y) > 2r+1$ forces the connection between the configurations $\omega_{B(0,r)}$ and $\omega_{B(y,r)}$ to be null. Hence, by the identity (3),

$$\begin{aligned} W_n(\omega) &= W_n(\omega_{B(0,r)} \omega_{B(y,r)}) \\ &= W_n(\omega_{B(0,r)}) W_n(\omega_{B(y,r)}) \\ &\leq W_n(\eta) W_n . \end{aligned}$$

Therefore, for any vertex $y \in \mathcal{R}(0, 2r+1, R(n)-r)$,

$$\mu_{a,b}(S_y^c) \geq 1 - |\mathcal{C}_r|^2 C_{2,r} W_n(\eta) W_n .$$

Hence, coupling this latter lower bound with (25) and Lemma 3.2, it follows that the conditional probability

$$\mu_{a,b}(\mathcal{A}_{r+1}(R(n), W_n) \mid I_0^\eta = 1)$$

is upper bounded by

$$(c_r W_n(\eta))^{-1} \left(1 - (1 - |\mathcal{C}_r|^2 C_{2,r} W_n(\eta) W_n)^{(2R(n)+1)^d} \right) . \quad (27)$$

Finally, the weights W_n and $W_n(\eta)$ and the product $R(n)^d W_n$ tend to 0. So, (27) is equivalent to

$$2^d c_r^{-1} |\mathcal{C}_r|^2 C_{2,r} R(n)^d W_n$$

and tends to 0 as n tends to infinity.

Let us turn to the second part of Theorem 5.3, i.e. statement (21). Some notations introduced in the previous section will be used here, starting with the set of vertices

$$T_n = \left\{ i(\rho(2r+1)+1), |i| = 0, \dots, \left\lfloor \frac{\rho R(n)}{\rho(2r+1)+1} \right\rfloor - 1 \right\}^d \cap B(0, R(n) - r)$$

(see Figure 2). Let $T_n^* = T_n \setminus \{0\}$ be the same set without the origin. If τ_n^* denotes the cardinality of T_n^* then there exists a constant $\tau^* > 0$ such that $\tau_n \geq \tau^* R(n)^d$. Let us denote by $\mathcal{T}_n^*$ the set

$$\mathcal{T}_n^* = \bigcup_{y \in T_n^*} B(y, r)$$

and by X_n^* the random variable

$$X_n^* = \sum_{y \in T_n^*} \sum_{\zeta \in \mathcal{C}_r(W_n)} I_y^\zeta.$$

The event $X_n^* > 0$ implies $\mathcal{A}_r(R(n), W_n)$. Hence, it suffices to prove that the conditional probability $\mu_{a,b}(X_n^* = 0 | I_0^\eta = 1)$ tends to 0. The end of the proof is very close to that of the upper bound of Theorem 4.1, so it will not be detailed.

The balls forming $\mathcal{T}_n^*$ are $\mathcal{V}$ -disjoint. So, by the Property 3.1, the probability $\mu_{a,b}(X_n^* = 0 \cap I_0^\eta = 1)$ can be expressed as

$$\mathbb{E}_{a,b} \left[\mu_{a,b}(I_0^\eta = 1 | \mathcal{F}(\delta B(0, r))) \times \prod_{y \in T_n^*} \mu_{a,b} \left(\bigcap_{\zeta \in \mathcal{C}_r(W_n)} I_y^\zeta = 0 \mid \mathcal{F}(\delta B(y, r)) \right) \right].$$

Thus, for any vertex $y \in T_n^*$, Lemma 3.2 provides

$$\mu_{a,b} \left(\bigcap_{\zeta \in \mathcal{C}_r(W_n)} I_y^\zeta = 0 \mid \mathcal{F}(\delta B(y, r)) \right) \leq 1 - c_r W_n.$$

As a consequence,

$$\mu_{a,b}(X_n^* = 0 \cap I_0^\eta = 1) \leq \mu_{a,b}(I_0^\eta = 1) (1 - c_r W_n)^{\tau_n^*}$$

and the conditional probability $\mu_{a,b}(X_n^* = 0 | I_0^\eta = 1)$ is now controlled:

$$\mu_{a,b}(X_n^* = 0 | I_0^\eta = 1) \leq e^{-c_r \tau^* R(n)^d W_n}$$

which tends to 0 since by hypothesis $R(n)^d W_n \rightarrow +\infty$. □

Although the proof of Theorem 5.4 seems to be more elaborated than that of Theorem 5.3, the ideas on which they are based are exactly the same.

Proof (of Theorem 5.4): Recall the definition of $\mathcal{H}_r(R(n), \ell(n), W_n)$:

$$\begin{aligned} \exists y, y' \in \mathcal{R}(0, 2r, R(n) - r), \exists \zeta, \zeta' \in \mathcal{D}_r(W_n) \text{ such that} \\ 2r < \text{dist}(y, y') \leq \ell(n) \text{ and } I_y^\zeta = I_{y'}^{\zeta'} = 1 . \end{aligned}$$

Let us start with the proof of (22). The first step consists in moving the occurrences of η , ζ and ζ' away from each other. For that purpose, let us introduce the 3 following events

$$\begin{aligned} \mathcal{H}^1 : \quad & \exists y \in V_n, \exists \zeta \in \mathcal{D}_r(W_n), \text{dist}(0, y) = 2r + 1 \text{ and } I_y^\zeta = 1 ; \\ \mathcal{H}^2 : \quad & \exists y, y' \in \mathcal{R}(0, 2r, R(n) - r), \exists \zeta, \zeta' \in \mathcal{D}_r(W_n) \text{ such that} \\ & \text{dist}(y, y') = 2r + 1 \text{ and } I_y^\zeta = I_{y'}^{\zeta'} = 1 ; \\ \mathcal{H}^3 : \quad & \exists y, y' \in \mathcal{R}(0, 2r + 1, R(n) - r), \exists \zeta, \zeta' \in \mathcal{D}_r(W_n) \text{ such that} \\ & 2r + 1 < \text{dist}(y, y') \leq \ell(n) \text{ and } I_y^\zeta = I_{y'}^{\zeta'} = 1 ; \end{aligned}$$

whose union contains $\mathcal{H}_r(R(n), \ell(n), W_n)$. Thanks to the hypothesis (H') , $\mathcal{H}^1$ is included in $\mathcal{A}_r(0, 3r + 1, +)$. So, Lemma 5.1 implies:

$$\lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{H}^1 | I_0^\eta = 1) = 0 .$$

Let y, y' be two vertices such that $\text{dist}(y, y') = 2r + 1$ and ζ, ζ' be two elements of $\mathcal{D}_r(W_n)$ occurring on the balls with radius r centered respectively at y and y' . The convexity of balls forces the connection between ζ_y and $\zeta'_{y'}$ to be smaller than $\mathcal{V}k(\zeta)$. Then, a bound for the weight of the configuration $\zeta_y \zeta'_{y'}$ is deduced:

$$\begin{aligned} W_n(\zeta_y \zeta'_{y'}) &= W_n(\zeta) W_n(\zeta') e^{4b(n) \text{conn}(\zeta_y, \zeta'_{y'})} \\ &\leq W_n^2 e^{2b(n) \mathcal{V}k_n} , \end{aligned}$$

where k_n is the index of W_n . In other words, the event $\mathcal{H}^2$ implies the existence in the ring $\mathcal{R}(0, r, R(n))$ of a local configuration with radius $3r + 1$ whose weight is smaller than $W_n^2 e^{2b(n) \mathcal{V}k_n}$. Now, by hypothesis

$$R(n)^d W_n^2 e^{2b(n) \mathcal{V}k_n}$$

tends to 0. So, it follows from Theorem 5.3:

$$\lim_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{H}^2 | I_0^\eta = 1) = 0 .$$

It remains to prove the same limit holds for the event $\mathcal{H}^3$. As at the end of the proof of Theorem 5.3, we are going to introduce increasing events in order to use the FKG

inequality. For any vertices y, y' , the event

$$S_{y,y'} = \left(\bigcup_{\zeta \in \mathcal{C}_r(W_n)} \{I_y^\zeta = 1\} \right) \cap \left(\bigcup_{\zeta' \in \mathcal{C}_r(W_n)} \{I_{y'}^{\zeta'} = 1\} \right) \cap \left(\bigcup_{\eta' \in \mathcal{C}_r(W_n(\eta))} \{I_0^{\eta'} = 1\} \right)$$

is increasing because $a(n) + \mathcal{V}b(n)$ is negative. Thus, the FKG inequality produces the following lower bound:

$$1 - \mu_{a,b}(\mathcal{H}^3 \cap \{I_0^\eta = 1\}) \geq \prod_{\substack{y \in \mathcal{R}(0, 2r+1, R(n)-r) \\ y' \in \mathcal{R}(y, 2r+1, \ell(n))}} \mu_{a,b}(S_{y,y'}^c) . \quad (28)$$

Let us pick $y \in \mathcal{R}(0, 2r+1, R(n)-r)$, $y' \in \mathcal{R}(y, 2r+1, \ell(n))$ and denote by $V_{y,y'}$ the union of the three balls $B(y, r)$, $B(y', r)$ and $B(0, r)$. Let $\mathcal{C}_{y,y'}$ be the set of configurations defined by:

$$\mathcal{C}_{y,y'} = \left\{ \omega \in \{-, +\}^{V_{y,y'}}, \begin{array}{l} W_n(\omega_{B(0,r)}) \leq W_n(\eta) \\ W_n(\omega_{B(y,r)}) \leq W_n \\ W_n(\omega_{B(y',r)}) \leq W_n \end{array} \right\} .$$

Then, using the inequality (7) of Lemma 3.2 (or rather its extension (26)), we bound the probability of $S_{y,y'}$. There exists a constant $C_{3,r} > 0$ such that

$$\begin{aligned} \mu_{a,b}(S_{y,y'}) &= \sum_{\omega \in \mathcal{C}_{y,y'}} \mu_{a,b}(I_{V_{y,y'}}^\omega = 1) \\ &\leq C_{3,r} \sum_{\omega \in \mathcal{C}_{y,y'}} W_n(\omega) . \end{aligned}$$

Now, vertices $0, y$ and y' are sufficiently far apart so that the weight of ω might write as the product of the weights of $\omega_{B(0,r)}$, $\omega_{B(y,r)}$ and $\omega_{B(y',r)}$. Then,

$$\mu_{a,b}(S_{y,y'}) \leq C_{3,r} |\mathcal{C}_r|^3 W_n(\eta) W_n^2$$

since $\omega \in \mathcal{C}_{y,y'}$ and the cardinality of $\mathcal{C}_{y,y'}$ is bounded by $|\mathcal{C}_r|^3$. Finally, coupling this latter inequality with (28) and Lemma 3.2, we bound the conditional probability $\mu_{a,b}(\mathcal{H}^3 | I_0^\eta = 1)$ by

$$(c_r W_n(\eta))^{-1} \left(1 - (1 - |\mathcal{C}_r|^3 C_{3,r} W_n(\eta) W_n^2)^{(2R(n)+1)^d (2\ell(n)+1)^d} \right) .$$

Since $W_n, W_n(\eta)$ and the product $\ell(n)^d R(n)^d W_n^2$ tend to 0, the above quantity is equivalent to

$$2^{2d} c_r^{-1} |\mathcal{C}_r|^3 C_{3,r} \ell(n)^d R(n)^d W_n^2$$

and tends to 0 as n tends to infinity. So does $\mu_{a,b}(\mathcal{H}^3 | I_0^\eta = 1)$.

Let us deal with the second part of Theorem 5.4, i.e. statement (23).

If the sequence $(\ell(n))_{n \in \mathbb{N}}$ were bounded, say by a constant ℓ_∞ , the event $\mathcal{H}_r(R(n), \ell(n), W_n)$ would correspond to the existence in the ring $\mathcal{R}(0, r, R(n))$ of a local configuration with

radius ℓ_∞ , say ζ , and whose weight would be larger than W_n^2 (by (3)). From then on, the limit

$$R(n)^d W_n(\zeta) \geq \frac{1}{\ell_\infty^d} R(n)^d \ell(n)^d W_n^2 \rightarrow +\infty$$

would imply (statement (21) of Theorem 5.3) a limit equal to 1 for the conditional probability $\mu_{a,b}(\mathcal{H}_r(R(n), \ell(n), W_n) | I_0^\eta = 1)$. So, from now on, we will assume the case where $\ell(n) \rightarrow +\infty$.

The second part of Theorem 5.4 needs one more time the set of vertices

$$T_n = \left\{ i(\rho(2r+1)+1), |i| = 0, \dots, \left\lfloor \frac{\rho R(n)}{\rho(2r+1)+1} \right\rfloor - 1 \right\}^d \cap B(0, R(n) - r)$$

(see Figure 2). Let $\phi_n = i_n(\rho(2r+1)+1)$, where

$$i_n = \left\lfloor \frac{\rho(\frac{4}{3}\ell(n)+1)+1}{\rho(2r+1)+1} \right\rfloor + 1,$$

and let us denote by $\mathcal{S}_n$ the sub-lattice of T_n defined by:

$$\mathcal{S}_n = \left\{ (\phi_n \mathbb{Z})^d \cap B\left(0, R(n) - \frac{2}{3}\ell(n)\right) \right\} \setminus \{0\}.$$

Any two balls $B(y, \frac{2}{3}\ell(n))$ and $B(y', \frac{2}{3}\ell(n))$, where $y, y' \in \mathcal{S}_n$, are $\mathcal{V}$ -disjoint since

$$\text{dist}(y, y') \geq \frac{\phi_n}{\rho} > \frac{4}{3}\ell(n) + 1.$$

Obviously, they are also included in the large ball $B(0, R(n))$. Let us pick a vertex y of $\mathcal{S}_n$. We denote by $\Lambda_1(y)$ and $\Lambda_2(y)$ the following subsets of T_n :

$$\Lambda_1(y) = \left\{ z \in T_n, \text{dist}(z, y) \leq \frac{1}{3}\ell(n) \right\}$$

and

$$\Lambda_2(y) = \left\{ z \in T_n, \frac{1}{3}\ell(n) < \text{dist}(z, y) \leq \frac{2}{3}\ell(n) \right\}.$$

Hence, two vertices $z \in \Lambda_1(y)$ and $z' \in \Lambda_2(y)$ satisfy $2r < \text{dist}(z, z') \leq \ell(n)$. As a consequence, the event

$$\mathcal{S} = \bigcup_{y \in \mathcal{S}_n} \bigcup_{\substack{z \in \Lambda_1(y) \\ z' \in \Lambda_2(y)}} \bigcup_{\zeta, \zeta' \in \mathcal{D}_r(W_n)} \left\{ I_z^\zeta = I_{z'}^{\zeta'} = 1 \right\}$$

implies $\mathcal{H}_r(R(n), \ell(n), W_n)$. So, it suffices to prove the conditional probability $\mu_{a,b}(\mathcal{S}^c | I_0^\eta = 1)$ tends to 0 as $n \rightarrow +\infty$.

The vertices of $\Lambda_1(y)$ are the elements of $((\rho(2r+1)+1)\mathbb{Z})^d$ whose distance (with respect

to $\|\cdot\|_\infty$) to y is smaller than $\frac{\ell}{3}\ell(n)$. Their number is of order $\ell(n)^d$. More generally, there exists a constant α which depends only on the parameters d, ρ, q and r , such that, for any vertex y ,

$$|\Lambda_i(y)| \geq \alpha \ell(n)^d, \quad i = 1, 2 \quad \text{and} \quad |\mathcal{S}_n| \geq \alpha \frac{R(n)^d}{\ell(n)^d}.$$

Moreover, the Markovian character of the measure $\mu_{a,b}$ (see Property 3.1) allows us to express the probability $\mu_{a,b}(\mathcal{S}^c \cap I_0^\eta = 1)$ as the following expectation:

$$\mathbb{E}_{a,b} \left[\mu_{a,b}(I_0^\eta = 1 \mid \mathcal{F}(\delta B(0, r))) \times \prod_{y \in \mathcal{S}_n} \prod_{\substack{z \in \Lambda_1(y) \\ z' \in \Lambda_2(y)}} \mu_{a,b} \left(\bigcap_{\zeta, \zeta' \in \mathcal{D}_r(W_n)} I_z^\zeta I_{z'}^{\zeta'} = 0 \mid \mathcal{F}(\delta V_{z, z'}) \right) \right],$$

where $V_{z, z'}$ is the union of the two balls $B(z, r)$ and $B(z', r)$. Let $y \in \mathcal{S}_n$ and $z \in \Lambda_1(y)$, $z' \in \Lambda_2(y)$. Let $\sigma \in \{-, +\}^{V_{z, z'}}$. Since the balls $B(z, r)$ and $B(z', r)$ $\mathcal{V}$ -disjoint,

$$\begin{aligned} \mu_{a,b} \left(\bigcap_{\zeta, \zeta' \in \mathcal{D}_r(W_n)} I_z^\zeta I_{z'}^{\zeta'} = 0 \mid \sigma \right) &= 1 - \sum_{\zeta, \zeta' \in \mathcal{D}_r(W_n)} \mu_{a,b}(I_z^\zeta = 0 \mid \sigma_{B(z, r)}) \mu_{a,b}(I_{z'}^{\zeta'} = 0 \mid \sigma_{B(z', r)}) \\ &\leq 1 - c_r^2 W_n^2. \end{aligned}$$

In conclusion,

$$\begin{aligned} \mu_{a,b}(\mathcal{S}^c \mid I_0^\eta = 1) &\leq (1 - c_r^2 W_n^2)^{\alpha^3 R(n)^d \ell(n)^d} \\ &\leq e^{-c_r^2 \alpha^3 R(n)^d \ell(n)^d W_n^2} \end{aligned}$$

which tends to 0 whenever $R(n)^d \ell(n)^d W_n^2$ tends to 0. $\square$

Proposition 5.5 is obtained using the results of the proofs of Lemma 5.1 and Theorem 5.3.

Proof (of Proposition 5.5): We are going to prove that the quantity $\mu_{a,b}(\mathcal{E}_n(\eta)^c \mid X_n(\eta) \geq 2)$ tends to 0 as $n \rightarrow +\infty$, where the event $\mathcal{E}_n(\eta)^c$ means

$$\forall C > 0, \exists x \neq y, I_x^\eta I_y^\eta = 1 \text{ and } \text{dist}(x, y) < Cn.$$

Thanks to the invariance translation of the graph G_n and for any constant $C > 0$,

$$\mu_{a,b}(\mathcal{E}_n(\eta)^c \mid X_n(\eta) \geq 2) \leq n^d \mu_{a,b}(\exists x \neq 0, I_x^\eta I_0^\eta = 1, \text{dist}(x, 0) \leq \lfloor Cn \rfloor \mid X_n(\eta) \geq 2).$$

The condition $X_n(\eta) \geq 2$ can be removed. Indeed, the random variable $X_n(\eta)$ is asymptotically Poissonian, under the hypothesis (H) (see [8]). So, there exist a constant $\varepsilon > 1$ and an integer $n(\varepsilon)$ such that the probability $\mu_{a,b}(X_n(\eta) \geq 2)$ is larger than ε^{-1} whenever $n \geq n(\varepsilon)$. Hence, for such value of n ,

$$\begin{aligned} \mu_{a,b}(\exists x \neq 0, I_x^\eta I_0^\eta = 1, \text{dist}(x, 0) \leq \lfloor Cn \rfloor \mid X_n(\eta) \geq 2) \\ \leq \varepsilon \mu_{a,b}(\exists x \neq 0, I_x^\eta I_0^\eta = 1, \text{dist}(x, 0) \leq \lfloor Cn \rfloor). \end{aligned}$$

In a first time, assume the centers 0 and x of balls with radius r on which the local configuration η occurs satisfy $\text{dist}(x, 0) \leq 2r + 1$. A configuration of $\{-, +\}^{B(0, R)}$, $R = 3r + 1$, fulfilling this event necessary belongs to

$$\mathcal{C}_R(\eta, +) = \{\zeta \in \mathcal{C}_R, \forall y \in B(0, r), \zeta(y) = \eta(y) \text{ and } k(\zeta) > k(\eta)\} .$$

Now, using the inequalities of the proof of Lemma 5.1, we get:

$$n^d \mu_{a,b}(\exists x \neq 0, I_x^\eta I_0^\eta = 1, \text{dist}(x, 0) \leq 2r + 1) \leq C_R |\mathcal{C}_R(\eta, +)| n^d W_n(\eta) e^{2a(n)}$$

which tends to 0 since C_R and $|\mathcal{C}_R(\eta, +)|$ are some constants, $n^d W_n(\eta) = \lambda$ and the magnetic field $a(n) \rightarrow -\infty$. So, the case where $\text{dist}(x, 0) \leq 2r + 1$ is negligible and

$$\begin{aligned} \limsup_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{E}_n(\eta)^c \mid X_n(\eta) \geq 2) \\ \leq \varepsilon \limsup_{n \rightarrow +\infty} n^d \mu_{a,b}(\exists x \neq 0, I_x^\eta I_0^\eta = 1, 2r + 1 < \text{dist}(x, 0) \leq \lfloor Cn \rfloor) , \end{aligned}$$

for any $C > 0$. Remark the event

$$\exists x \neq 0, I_x^\eta I_0^\eta = 1, 2r + 1 < \text{dist}(x, 0) \leq \lfloor Cn \rfloor$$

is included in $\mathcal{A}_{r+1}(0, \lfloor Cn \rfloor, W_n(\eta)) \cap \{I_0^\eta = 1\}$. In the proof of Theorem 5.3, it has been proved that

$$\mu_{a,b}(\mathcal{A}_{r+1}(0, \lfloor Cn \rfloor, W_n(\eta)) \cap \{I_0^\eta = 1\}) = \mathcal{O}(\lfloor Cn \rfloor^d W_n(\eta)^2) .$$

Finally, since the product $n^d W_n(\eta)$ is constant, we get:

$$n^d \mu_{a,b}(\exists x \neq 0, I_x^\eta I_0^\eta = 1, 2r + 1 < \text{dist}(x, 0) \leq \lfloor Cn \rfloor) = \mathcal{O}(C^d) .$$

Take $C \searrow 0$ and the desired result follows:

$$\limsup_{n \rightarrow +\infty} \mu_{a,b}(\mathcal{E}_n(\eta)^c \mid X_n(\eta) \geq 2) = 0 .$$

□

6 Ubiquity of local configurations

The goal of this section is to give a criterion ensuring that a given local configuration occurs everywhere in the graph. This criterion is presented in Proposition 6.2 through the use of an appropriate Markovian measure. So, the first step is to build a probabilistic model from the Ising model of Definition 1.1.

A simple way to cover the set of vertices V_n by balls consists in using the L_∞ norm. In this case, replacing the radius r with ρr , we can assume that $\rho = 1$. So, in this section,

the neighborhood structure of each vertex is $\mathcal{V}(x) = \{y \in V_n, \|y - x\|_\infty = 1\}$. Hence, the graph distance $dist$ and the L_∞ norm define the same sets; the ball $B(x, r)$ is equal to the hypercube with center x and radius r .

For all integer n , let us denote by $v(n)$ the largest integer m such that 2^m divides n and by $R(n)$ the integer satisfying the relation

$$\frac{n}{2^{v(n)}} = 2R(n) + 1 .$$

A function $g : \mathbb{N}^* \rightarrow \mathbb{N}^*$ satisfying for all integer n the inequalities $v(g(n+1)) \geq v(g(n))$ and $R(g(n+1)) \geq R(g(n))$, and tending to infinity as $n \rightarrow +\infty$ will be said *adequate*. In particular, an adequate function is nondecreasing. The functions recursively defined by

$$\begin{cases} g(n+1) = 2^{v(g(n))+1} (2(R(g(n)) + 1) + 1), \forall n \geq 1 \\ g(1) \in \mathbb{N}^* \end{cases}$$

provide examples of adequate functions, since $v(g(n+1))$ and $R(g(n+1))$ are respectively equal to $v(g(n)) + 1$ and $R(g(n)) + 1$. For instance, if $g(1) = 1$ then $g(2) = 2^1 \times 3 = 6$, $g(3) = 2^2 \times 5 = 20$, $g(4) = 2^3 \times 7 = 56$...

In conclusion, replacing n with $g(n)$ where g is an adequate function, we will assume that there exists a nondecreasing, integer valued sequence $(R(n))_{n \geq 1}$ such that the sequence

$$\left(\frac{n}{2R(n) + 1} \right)_{n \geq 1}$$

is nondecreasing and integer valued too.

Now, let us denote by $\tilde{V}_n$ the following subset of V_n :

$$\tilde{V}_n = \left\{ i(2R(n) + 1), i = 0, \dots, \frac{n}{2R(n) + 1} - 1 \right\}^d .$$

The set of balls $\{B(x, R(n)), x \in \tilde{V}_n\}$ is a partition of V_n . The edge set $\tilde{E}_n$ is specified by defining the neighborhood $\tilde{\mathcal{V}}(x)$ of each vertex $x \in \tilde{V}_n$:

$$\tilde{\mathcal{V}}(x) = \{y \in \tilde{V}_n, \|y - x\|_\infty = 2R(n) + 1\} . \quad (29)$$

By analogy with the previous sections, we denote by $\delta\tilde{V}$ the neighborhood of $\tilde{V} \subset \tilde{V}_n$ corresponding to (29). Hence, an undirected graph structure $\tilde{G}_n = (\tilde{V}_n, \tilde{E}_n)$ with periodic boundary conditions is defined. The *size* of $\tilde{G}_n$ is the ratio n divided by $2R(n) + 1$. Furthermore, remark the graph $\tilde{G}_n$ retains the translation invariance property of G_n .

Let η be a local configuration with radius r . We associate to η a function f from $\mathcal{X}_n$ into $\tilde{\mathcal{X}}_n = \{-, +\}^{\tilde{V}_n}$ defined for $\sigma \in \mathcal{X}_n$ and $x \in \tilde{V}_n$ by;

$$f(\sigma)(x) = + \iff \exists y \in B(x, R(n)), I_y^\eta(\sigma) = 1 .$$

In other words, the vertex $x \in \tilde{V}_n$ is positive for the configuration $f(\sigma) \in \tilde{\mathcal{X}}_n$ if and only if a copy of η occurs in the ball $B(x, R(n) + r)$ (under σ). Remark the function f is onto $\tilde{\mathcal{X}}_n$. In particular, for $\tilde{V} \subset \tilde{V}_n$, the values taken by a configuration $\sigma \in \mathcal{X}_n$ on the set

$$\Sigma = \bigcup_{y \in \tilde{V}} B(y, R(n) + r)$$

specify completely the configuration $f(\sigma)$ on $\tilde{V}$. Besides, f is not an injective function; for $\zeta \in \{-, +\}^{\tilde{V}}$, $f^{-1}(\zeta)$ represents a subset of $\{-, +\}^{\Sigma}$.

Let $\mathcal{F}(\tilde{V})$ be the σ -algebra generated by the configurations of $\{-, +\}^{\tilde{V}}$. Then, $f^{-1}(\mathcal{F}(\tilde{V}))$ is still a σ -algebra, generated by the sets $f^{-1}(\zeta)$, $\zeta \in \{-, +\}^{\tilde{V}}$, and is coarser than $\mathcal{F}(\Sigma)$:

$$f^{-1}(\mathcal{F}(\tilde{V})) \subsetneq \mathcal{F}(\Sigma) .$$

Thus, let us endow the set of configurations $\tilde{\mathcal{X}}_n$ with the measure $\tilde{\mu}_{a,b}$ defined by:

$$\forall \zeta \in \tilde{\mathcal{X}}_n, \tilde{\mu}_{a,b}(\zeta) = \mu_{a,b}(f^{-1}(\zeta)) . \quad (30)$$

Expectations relative to $\tilde{\mu}_{a,b}$ will be denoted by $\tilde{E}_{a,b}$.

Property 6.1 links the random variable $\tilde{\mu}_{a,b}(\cdot | \mathcal{F}(\tilde{V}))$ to $\mu_{a,b}(\cdot | f^{-1}(\mathcal{F}(\tilde{V})))$ and states the Markovian character of $\tilde{\mu}_{a,b}$. The identity (31) is completely based on the definition of the measure $\tilde{\mu}_{a,b}$. It holds whatever the function f . Relation (32) derives from the Markovian character of the Gibbs measure $\mu_{a,b}$ and the use of the L_∞ norm in the construction of the graph $\tilde{G}_n$. Property 6.1 will be proved at the end of the section.

Property 6.1 *Let $\tilde{U}, \tilde{V}$ be two subsets of $\tilde{V}_n$. Then, for all event $A \in \mathcal{F}(\tilde{U})$,*

$$\tilde{\mu}_{a,b} \left(A | \mathcal{F}(\tilde{V}) \right) \circ f = \mu_{a,b} \left(f^{-1}(A) | f^{-1}(\mathcal{F}(\tilde{V})) \right) , \quad (31)$$

where $\circ$ denotes the composition relation. Moreover, if $\tilde{U} \cap \tilde{V} = \emptyset$ and $\delta\tilde{U} \subset \tilde{V}$ then, for all event $A \in \mathcal{F}(\tilde{U})$,

$$\tilde{\mu}_{a,b} \left(A | \mathcal{F}(\tilde{V}) \right) = \tilde{\mu}_{a,b} \left(A | \mathcal{F}(\delta\tilde{U}) \right) . \quad (32)$$

The rest of this section is devoted to the study of the asymptotic behavior of the probability measure $\tilde{\mu}_{a,b}$. As n tends to infinity, the two following sequences

$$(R(n))_{n \geq 1} \quad \text{and} \quad \left(\frac{n}{2R(n) + 1} \right)_{n \geq 1}$$

cannot be simultaneously bounded. Then, three alternatives are conceivable; either the graph G_n is divided into a small number of large balls, or into a large number of small balls, or into a large number of large balls.

Proposition 6.2 gives sufficient conditions describing the asymptotic behavior of $\tilde{\mu}_{a,b}$. Relations (33) and (34) are a rewriting of results already known; the second one means at

least one copy of the local configuration η can be found somewhere in the graph, with probability tending to 1 (for $\mu_{a,b}$), whenever $W_n(\eta)$ is larger than n^{-d} . Besides, recall that any given ball $B(x, R(n) + r)$ contains a copy of η whenever $W_n(\eta)$ is larger than $R(n)^{-d}$ (Theorem 4.1). But, so as to every ball $B(x, R(n) + r)$, $x \in \tilde{V}_n$, contains a copy of η (this is that we call *ubiquity* of the local configuration η) a stronger condition is given by Proposition 6.2; $W_n(\eta)$ must be larger than $R(n)^{-d} \ln(n/R(n))$.

For that purpose, let ζ_- and ζ_+ be the two configurations of $\tilde{\mathcal{X}}_n$ whose vertices are all negative and all positive.

Proposition 6.2 *Let us consider a local configuration $\eta \in \mathcal{C}_r$ and potentials $a(n) < 0$ and $b(n) \geq 0$ such that $a(n) + 2\mathcal{V}b(n) \leq 0$.*

$$\text{If } \lim_{n \rightarrow +\infty} n^d W_n(\eta) = 0 \text{ then } \lim_{n \rightarrow +\infty} \tilde{\mu}_{a,b}(\zeta_-) = 1 . \quad (33)$$

$$\text{If } \lim_{n \rightarrow +\infty} n^d W_n(\eta) = +\infty \text{ then } \lim_{n \rightarrow +\infty} \tilde{\mu}_{a,b}(\zeta_-) = 0 . \quad (34)$$

$$\text{If } \lim_{n \rightarrow +\infty} R(n)^d \ln \left(\frac{n}{R(n)} \right)^{-1} W_n(\eta) = +\infty \text{ then } \lim_{n \rightarrow +\infty} \tilde{\mu}_{a,b}(\zeta_+) = 1 . \quad (35)$$

Relations (33) and (35) respectively say that $\tilde{\mu}_{a,b}$ converges weakly to the Dirac measures associated to the configurations ζ_- and ζ_+ .

The probability (for $\tilde{\mu}_{a,b}$) for a given vertex $x \in \tilde{V}_n$ to be positive is equal to the probability (for $\mu_{a,b}$) that the local configuration η occurs somewhere in the ball $B(x, R(n) + r)$; see relation (36) below. Now, this quantity has been studied and bounded in Section 4. The proof of Proposition 6.2 immediatly derives from this remark.

Proof (of Proposition 6.2): For $x \in \tilde{V}_n$, let $\tilde{I}_x^+$ be the indicator function defined on $\tilde{\mathcal{X}}_n$ as follows: $\tilde{I}_x^+(\zeta)$ is 1 if the vertex x is positive under $\zeta \in \tilde{\mathcal{X}}_n$, i.e. $\zeta(x) = 1$, and 0 otherwise. Then,

$$\begin{aligned} \tilde{\mu}_{a,b} \left(\exists x \in \tilde{V}_n, \tilde{I}_x^+ = 1 \right) &= \mu_{a,b} (\exists x \in V_n, I_x^\eta = 1) \\ &= \mu_{a,b} (X_n(\eta) > 0) , \end{aligned}$$

where $X_n(\eta)$ represents the number of copies of η occurring in G_n . In [8], it has been proved that $\mu_{a,b}(X_n(\eta) > 0)$ tends to 0 (resp. 1) whenever $n^d W_n(\eta)$ tends to 0 (resp. $+\infty$). Relations (33) and (34) follow.

In order to obtain (35), we are going to prove that the probability of the opposite event $\exists x \in \tilde{V}_n, \tilde{I}_x^+ = 0$ tends to 0. Due to the translation invariance of $\tilde{G}_n$, the random indicators $\tilde{I}_x^+$ have the same distribution. So,

$$\tilde{\mu}_{a,b} \left(\exists x \in \tilde{V}_n, \tilde{I}_x^+ = 0 \right) \leq \left(\frac{n}{R(n)} \right)^d \tilde{\mu}_{a,b} \left(\tilde{I}_0^+ = 0 \right) .$$

Furthermore, following the inequalities of the proof of the upper bound of (11), we bound the probability of the event $\tilde{I}_0^+ = 0$:

$$\begin{aligned}\tilde{\mu}_{a,b} \left(\tilde{I}_0^+ = 0 \right) &= \mu_{a,b} \left(\tilde{I}_0^+ \circ f = 0 \right) \\ &= \mu_{a,b} \left(\sum_{y \in B(0, R(n))} I_y^\eta = 0 \right) \\ &\leq \exp \left(-\tau c_r R(n)^d W_n(\eta) \right) ,\end{aligned}\tag{36}$$

where τ and c_r are positive constant. Finally, (35) is deduced from

$$\tilde{\mu}_{a,b} \left(\exists x \in \tilde{V}_n, \tilde{I}_x^+ = 0 \right) \leq \exp \left(-R(n)^d W_n(\eta) \left(\tau c_r - \frac{d \ln(n/R(n))}{R(n)^d W_n(\eta)} \right) \right) .$$

□

Proof (of Property 6.1): First, note that any event $A \in \mathcal{F}(\tilde{U})$ can be written as a disjoint union of configurations of $\{-, +\}^{\tilde{U}}$. So, it suffices to prove the identities (31) and (32) for $A = \{\zeta\}$, $\zeta \in \{-, +\}^{\tilde{U}}$.

Let us pick such a configuration $\zeta \in \{-, +\}^{\tilde{U}}$. The set $\{f^{-1}(\zeta'), \zeta' \in \{-, +\}^{\tilde{V}}\}$ is a π -system which generates the σ -algebra $f^{-1}(\mathcal{F}(\tilde{V}))$. Hence, it is enough to prove

$$\mathbb{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \circ f \mathbb{1}_{f^{-1}(\zeta')} \right] = \mathbb{E}_{a,b} \left[\mathbb{1}_{f^{-1}(\zeta)} \mathbb{1}_{f^{-1}(\zeta')} \right], \quad \forall \zeta' \in \{-, +\}^{\tilde{V}} \tag{37}$$

and

$$\mathbb{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \circ f \right] = \mathbb{E}_{a,b} \left[\mathbb{1}_{f^{-1}(\zeta)} \right] \tag{38}$$

(see for example [23] p.84). Let us start with relation (37). For a configuration ζ' belonging to $\{-, +\}^{\tilde{V}}$,

$$\begin{aligned}\mathbb{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \circ f \mathbb{1}_{f^{-1}(\zeta')} \right] &= \mathbb{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \circ f \mathbb{1}_{\zeta'} \circ f \right] \\ &= \tilde{\mathbb{E}}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \mathbb{1}_{\zeta'} \right] \\ &= \tilde{\mathbb{E}}_{a,b} \left[\mathbb{1}_{\zeta} \mathbb{1}_{\zeta'} \right] \\ &= \mathbb{E}_{a,b} \left[\mathbb{1}_{f^{-1}(\zeta)} \mathbb{1}_{f^{-1}(\zeta')} \right] .\end{aligned}$$

Relation (38) is treated in the same way:

$$\begin{aligned}\mathbb{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \circ f \right] &= \tilde{\mathbb{E}}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \right] \\ &= \tilde{\mathbb{E}}_{a,b} \left[\mathbb{1}_{\zeta} \right] \\ &= \mathbb{E}_{a,b} \left[\mathbb{1}_{f^{-1}(\zeta)} \right] .\end{aligned}$$

Now, let us prove (32) with $A = \{\zeta\}$, $\zeta \in \{-, +\}^{\tilde{U}}$. The set $\{\zeta', \zeta' \in \{-, +\}^{\delta\tilde{U}}\}$ is a π -system which generates the σ -algebra $\mathcal{F}(\delta\tilde{U})$. So, since the random variables $\tilde{\mu}_{a,b}(\zeta|\mathcal{F}(\tilde{V}))$ and $\tilde{\mu}_{a,b}(\zeta|\mathcal{F}(\delta\tilde{U}))$ have the same expectation (equal to $\tilde{\mu}_{a,b}(\zeta)$), it suffices to prove that, for any $\zeta' \in \{-, +\}^{\delta\tilde{U}}$,

$$\tilde{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta | \mathcal{F}(\tilde{V}) \right) \mathbb{1}_{\zeta'} \right] = \tilde{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta | \mathcal{F}(\delta\tilde{U}) \right) \mathbb{1}_{\zeta'} \right]. \quad (39)$$

Let ζ' be a configuration of $\{-, +\}^{\delta\tilde{U}}$. Firstly, relation (31) allows us to express the above expectations according to the measure $\mu_{a,b}$:

$$\begin{aligned} \tilde{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta | \mathcal{F}(\tilde{V}) \right) \mathbb{1}_{\zeta'} \right] &= E_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta | \mathcal{F}(\tilde{V}) \right) \circ f \mathbb{1}_{\zeta'} \circ f \right] \\ &= E_{a,b} \left[\mu_{a,b} \left(f^{-1}(\zeta) | f^{-1}(\mathcal{F}(\tilde{V})) \right) \mathbb{1}_{f^{-1}(\zeta')} \right]. \end{aligned}$$

Thus, let us denote by Σ_1 and Σ_2 the following sets:

$$\Sigma_1 = \bigcup_{y \in \delta\tilde{U}} B(y, R(n) + r) \quad \text{and} \quad \Sigma_2 = \left(\bigcup_{y \in \tilde{V}} B(y, R(n) + r) \right) \setminus \Sigma_1.$$

Since $f^{-1}(\zeta')$ is a subset of Σ_1 , we can write:

$$\begin{aligned} \tilde{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta | \mathcal{F}(\tilde{V}) \right) \mathbb{1}_{\zeta'} \right] &= \sum_{\substack{\omega \in \{-, +\}^{\Sigma_1} \\ f(\omega) = \zeta'}} E_{a,b} \left[\mu_{a,b} \left(f^{-1}(\zeta) | f^{-1}(\mathcal{F}(\tilde{V})) \right) \mathbb{1}_{\omega} \right] \\ &= \sum_{\substack{\omega \in \{-, +\}^{\Sigma_1} \\ f(\omega) = \zeta'}} \sum_{\omega' \in \{-, +\}^{\Sigma_2}} E_{a,b} \left[\mu_{a,b} \left(f^{-1}(\zeta) | f^{-1}(\mathcal{F}(\tilde{V})) \right) \mathbb{1}_{\omega\omega'} \right], \end{aligned}$$

where $\omega\omega'$ is a configuration of $\{-, +\}^{\Sigma_1 \cup \Sigma_2}$. Now, the variable $\mu_{a,b}(f^{-1}(\zeta)|f^{-1}(\mathcal{F}(\tilde{V})))$ only depends on the vertices of $\Sigma_1 \cup \Sigma_2$. Hence,

$$E_{a,b} \left[\mu_{a,b} \left(f^{-1}(\zeta) | f^{-1}(\mathcal{F}(\tilde{V})) \right) \mathbb{1}_{\omega\omega'} \right] = \mu_{a,b} \left(f^{-1}(\zeta) | \omega\omega' \right) \mu_{a,b}(\omega\omega').$$

Furthermore, the configurations belonging to $f^{-1}(\zeta)$ only depend on the vertices of balls $B(y, R(n) + r)$, $y \in \tilde{U}$, and by construction of the graph $\tilde{G}_n$ (and the use of the L_∞ norm) the following inclusion holds:

$$\delta \left(\bigcup_{y \in \tilde{U}} B(y, R(n) + r) \right) \subset \Sigma_1.$$

So, the Markovian character of the measure $\mu_{a,b}$ applies. The conditional probability $\mu_{a,b}(f^{-1}(\zeta)|\omega\omega')$ can be reduced to $\mu_{a,b}(f^{-1}(\zeta)|\omega)$ (see [21] p.7). Combining the previous

equalities, it follows:

$$\begin{aligned}
\tilde{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \mathbb{1}_{\zeta'} \right] &= \sum_{\substack{\omega \in \{-,+\}^{\Sigma_1} \\ f(\omega) = \zeta'}} \mu_{a,b} \left(f^{-1}(\zeta) \mid \omega \right) \sum_{\omega' \in \{-,+\}^{\Sigma_2}} \mu_{a,b}(\omega \omega') \\
&= \sum_{\substack{\omega \in \{-,+\}^{\Sigma_1} \\ f(\omega) = \zeta'}} \mu_{a,b} \left(f^{-1}(\zeta) \mid \omega \right) \mu_{a,b}(\omega) \\
&= \sum_{\substack{\omega \in \{-,+\}^{\Sigma_1} \\ f(\omega) = \zeta'}} E_{a,b} \left[\mu_{a,b} \left(f^{-1}(\zeta) \mid f^{-1}(\mathcal{F}(\delta \tilde{U})) \right) \mathbb{1}_{\omega} \right] \\
&= E_{a,b} \left[\mu_{a,b} \left(f^{-1}(\zeta) \mid f^{-1}(\mathcal{F}(\delta \tilde{U})) \right) \mathbb{1}_{f^{-1}(\zeta')} \right].
\end{aligned}$$

Finally, using a second time the relation (31), we get the desired identity:

$$\begin{aligned}
\tilde{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\tilde{V}) \right) \mathbb{1}_{\zeta'} \right] &= E_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\delta \tilde{U}) \right) \circ f \mathbb{1}_{\zeta'} \circ f \right] \\
&= \tilde{E}_{a,b} \left[\tilde{\mu}_{a,b} \left(\zeta \mid \mathcal{F}(\delta \tilde{U}) \right) \mathbb{1}_{\zeta'} \right].
\end{aligned}$$

□

References

- [1] M. Abadi, J.-R. Chazottes, F. Redig, and E. Verbitskiy. Exponential distribution for the occurrence of rare patterns in Gibbsian random fields. *Comm. Math. Phys.*, 246(2):269–294, 2004.
- [2] M. Abadi and A. Galves. Inequalities for the occurrence times of rare events in mixing processes. The state of the art. *Markov Process. Related Fields*, 7(1):97–112, 2001. Inhomogeneous random systems (Cergy-Pontoise, 2000).
- [3] A.D. Barbour, L. Holst, and S. Janson. *Poisson approximation*. Oxford University Press, New York, 1992.
- [4] B. Bollobás. *Random Graphs*. Academic Press, London, 1985.
- [5] R. Boppona and J. Spencer. A useful elementary correlation inequality. *J. Combin. Theory Ser. A*, 50(2):305–307, 1989.
- [6] J.-R. Chazottes and F. Redig. Occurrence, repetition and matching of patterns in the low-temperature Ising model. *J. Stat. Phys.*, 121(3-4), 2005.
- [7] L.H.Y. Chen. Poisson approximation for dependent trials. *Ann. Probability*, 3(3):534–545, 1975.

- [8] D. Coupier. Two sufficient conditions for Poisson approximations in the ferromagnetic the Ising model. *Submitted*,
<http://math.univ-lille1.fr/~coupier/papier/2sufficient.pdf>.
- [9] D. Coupier. *Asymptotique des propriétés locales pour le modèle d'Ising et applications*. PhD thesis, Université Paris 5, 2005.
- [10] D. Coupier. Poisson approximations for the Ising model. *J. Stat. Phys.*, 123(2):473–495, 2006.
- [11] D. Coupier, P. Doukhan, and B. Ycart. Zero-one laws for binary random fields. *ALEA Lat. Am. J. Probab. Math. Stat.*, 2:157–175 (electronic), 2006.
- [12] P. Erdős and A. Rényi. On the evolution of random graphs. *Mat. Kuttató. Int. Közl.*, 5:17–60, 1960.
- [13] R. Fernández, P.A. Ferrari, and N.L. Garcia. Loss network representation of Peierls contours. *Ann. Probab.*, 29(2):902–937, 2001.
- [14] P.A. Ferrari and P. Picco. Poisson approximation for large-contours in low-temperature Ising models. *Physica A: Statistical Mechanics and its Applications*, 279(1-4):303–311, 2000.
- [15] C.M. Fortuin, P.W. Kasteleyn, and J. Ginibre. Correlation inequalities on some partially ordered sets. *Comm. Math. Phys.*, 22:89–103, 1971.
- [16] A. Ganesh, B.M. Hambly, N. O'Connell, D. Stark, and P.J. Upton. Poissonian behavior of Ising spin systems in an external field. *J. Statist. Phys.*, 99(1-2):613–626, 2000.
- [17] H.O. Georgii. *Gibbs measures and phase transitions*. de Gruyter, Berlin, 1988.
- [18] G. Grimmett. *Percolation*. Springer-Verlag, New York, 1989.
- [19] S. Janson. Poisson approximation for large deviations. *Random Structures Algorithms*, 1(2):221–229, 1990.
- [20] S. Janson, T. Łuczak, and A. Ruciński. An exponential bound for the probability of nonexistence of a specified subgraph in a random graph. In *Random graphs '87 (Poznań, 1987)*, pages 73–87. Wiley, Chichester, 1990.
- [21] V.A. Malyshev and R.A. Minlos. *Gibbs Random Fields, Cluster Expansions*. Kluwer Academic Publishers, 1991.
- [22] J. Spencer. *The Strange Logic on Random Graphs*, volume 22 of *Algorithms and Combinatorics*. Springer, 2001.
- [23] D. Williams. *Probability with Martingales*. Cambridge University Press, 1991.